arXiv:0910.0358v3 [math.DG] 1 Sep 2010

TORUS FIBRATIONS AND LOCALIZATION OF INDEX II
- LOCAL INDEX FOR ACYCLIC COMPATIBLE SYSTEM -

HAJIME FUJITA[1], MIKIO FURUTA[2], AND TAKAHIKO YOSHIDA[3]

Abstract. We give a framework of localization for the index of a Dirac-type operator on an open manifold. Suppose the open manifold has a compact subset whose complement is covered by a family of finitely many open subsets, each of which has a structure of the total space of a torus bundle. Under some compatibility and acyclicity conditions we show that the index of the Dirac-type operator is localized on the compact set.

Contents

1. Introduction 2
2. Compatible fibration and acyclic compatible system 3
2.1. Compatible fibration 3
2.2. Acyclic compatible system 7
2.3. Example from torus action 11
3. An index theory for complete Riemannian manifolds 16
3.1. Formulation of index on complete manifolds 16
3.2. Partial integration 17
3.3. Min-max principle 19
3.4. Deformation invariance of index 21
3.5. Gluing formula 22
3.6. Product formula 23
4. Local index 24
4.1. Vanishing theorem 26
4.2. Definition of the local index - Generalized cylindrical end case 29
4.3. Definition of the local index - general case 30
5. Product formula of local indices 31
5.1. Product of compatible fibrations 31
5.2. Product of acyclic compatible systems 32
5.3. Product formula 34
6. Four-dimensional case 35
6.1. Local indices for elliptic singularities 35
6.2. Application to locally toric Lagrangian fibrations 38
Appendix A. Proof of Proposition 2.15 45
Appendix B. A sufficient condition for Assumption 2.12 (2) 47

2000 *Mathematics Subject Classification.* Primary 19K56; Secondary 58J05, 53D50.

Key words and phrases. index theory, localization.

[1] Partly supported by Grant-in-Aid for Young Scientists (Start-up) 21840045.

[2] Partly supported by Grant-in-Aid for Scientific Research (A) 19204003 and Grant-in-Aid for Scientific Research (B) 19340015.

[3] Partly supported by Grant-in-Aid for Young Scientists (B) 20740029, and Fujyukai Foundation.

Appendix C. Proof of Lemma 2.30 47
Appendix D. Proof of Lemma 3.10 48
References 49

1. Introduction

This paper is the second of the series concerning a localization of the index of elliptic operators.

For a linear elliptic operator on a closed manifold, its Fredholm index is sometimes determined by the information on a specific subset under appropriate geometric conditions. Such a phenomenon is called *localization of index*. A typical example is Hopf's theorem identifying the index of the de Rham operator with the number of zeros of a vector field counted with sign and multiplicity. In this case the geometric condition is given by the vector field, and the index is localized around the zeros of the vector field. Another typical example is Atiyah-Segal's Lefschetz formula for the equivariant index for a torus action, when the geometric condition is given by the torus action. The index is localized around its fixed point set, and the localization is understood in terms of an algebraic localization of equivariant K-group. In particular when the manifold is symplectic and the elliptic operator is a Dirac-type operator, the localization is extensively investigated using the relation between the algebraic localizations in equivariant K-theory and that in equivariant ordinary cohomology theory.

In the previous paper [4], we investigated the case of a closed symplectic manifold equipped with a prequantizing line bundle and the structure of a Lagrangian fibration, and described a localization of the index of a Dirac-type operator, twisted by the prequantizing line bundle, on the subset consisting of Bohr-Sommerfeld fibers and singular fibers. A novel feature of our method is that we do not use a global group action but use only the structure of a torus bundle on an open subset of the manifold.

In the present paper we generalize our method to deal with the case when we do not have a global torus bundle on the open subset, but we just have the structure of a torus bundle on a neighborhood of each points, which gives a family of torus bundles satisfying some compatibility condition. The various torus bundles may have tori of various dimensions as their fibers. This generalization enables us to describe the localization phenomenon more precisely. Even for the case in the previous paper, we could replace the subset on which the index is localized with a smaller subset. A typical example of our generalization is the localization for the Riemann-Roch number of toric manifolds, for which we would need an orbifold version of our formulation. Moreover we can deal with some prequantized singular Lagrangian fibration without global toric action (Section 6). In our subsequent paper we will use the localization to give an approach to V. Guillemin and S. Sternberg's conjecture concerning "quantization commutes with reduction" in the case of torus actions. Though our motivating example is the Riemann-Roch number of a symplectic manifold, the localization of the index is formulated for more general cases. In fact we first establish a general framework to formulate the index of elliptic operators on complete manifolds (Section 3). This section is independent of the other sections and the framework might itself be interesting.

The mechanism of our localization is explained as a version of Witten's deformation, where the potential term itself is a first order differential operator. Our geometric input data is a family of torus bundles. Roughly speaking we deform the operator like an *adiabatic limit* shrinking the various fiber directions at the same time in a compatible manner. The *potential term* corresponds to some average of the de Rham operators along the various fiber directions.

Formally our localization is formulated as a property for the *index* of the elliptic operator on an open manifold: let D be an elliptic operator on a (possibly non-compact) manifold X, and V is an open subset of X whose complement $X \smallsetminus V$ is compact. Suppose V has a certain geometric structure s, by which we can modify D to construct a Fredholm operator. The index of the Fredholm operator depends on the data (X, V, s, D). Suppose the index satisfies the following properties. Firstly the index is deformation invariant. Secondly if X' is an open subset of X containing $X \smallsetminus V$, and hence $X' \smallsetminus V$ is compact. Let D' be the restriction of D on X'. We assume that the structure s has its *restriction* s' on $V' = X' \cap V$. Then we have the index of the Fredholm operator constructed from the data (X', V', s', D'). The required excision property is the equality between the two indices. We will construct Fredholm operators which satisfy the above type of excision property. The structure s on V is not extended on the whole X. In this sense $X \smallsetminus V$ is regarded as the *singular locus* of the structure. The index is *localized* on the singular locus $X \smallsetminus V$, and we call it the *local index* of the data (X, V, s, D). When $X \smallsetminus V$ is of the form of the disjoint union of finitely many compact subsets, the *localized* index is equal to the sum of the contributions from the compact subsets.

Our first main result is the construction of the local index when the structure s is the *strongly acyclic compatible system* defined in Section 2. Our second main result is a few basic properties of the local index, in particular a *product formula* of the local index.

The organization of this paper is as follows. In Section 2 we define the notion of a strongly acyclic compatible system, which we use as the geometric structure s in the above explanation. In Section 3 we give a formulation of the index of elliptic operators on complete Riemannian manifolds. This formulation is a generalization of the one given in Section 5 of [6]. This section is independent of the other sections. In Section 4 we define the index of elliptic operators using the framework of Section 3 under the assumption that a strongly acyclic compatible system is given on an end of the base manifold. In Section 5 we show a product formula for the index defined in Section 4. In Section 6 we give an example of our formulation using some 4-dimensional Lagrangian fibration with singular fibers. In Appendix we give proofs of the technical lemmas used in the main part.

Acknowledgments. The authors would like to thank Yukio Kametani, Shinichiroh Matsuo, Nobuhiro Nakamura and Hirohumi Sasahira. Throughout the seminar with them the authors could simplify the proof of the key proposition (Proposition 4.10). The second author is grateful to the hospitality of MIT and University of Minnesota.

2. Compatible fibration and acyclic compatible system

In this paper we work in smooth category. In particular, every manifold is a smooth manifold and every fiber bundle is a smooth fiber bundle.

2.1. Compatible fibration.

Let M be a manifold.

Definition 2.1. A *compatible fibration on M* is a collection of data

- $\{\pi_\alpha \colon V_\alpha \to U_\alpha \mid \alpha \in A\}$,
- $\{\pi_{\alpha\beta} \colon V_\alpha \cap V_\beta \to U_{\alpha\beta} \mid \alpha, \beta \in A$ such that $V_\alpha \cap V_\beta \neq \emptyset$ and $\alpha \neq \beta\}$, and

- $\{p^\alpha_{\alpha\beta} \colon U_{\alpha\beta} \to \pi_\alpha(V_\alpha \cap V_\beta) \mid \alpha, \beta \in A$ such that $V_\alpha \cap V_\beta \neq \emptyset$ and $\alpha \neq \beta\}$

satisfying the following properties.

(1) $\{V_\alpha\}_{\alpha \in A}$ is an open covering of M.

(2) For each $\alpha \in A$, U_α is a manifold and $\pi_\alpha \colon V_\alpha \to U_\alpha$ is a fiber bundle whose fiber is a closed manifold.

(3) For each α and β, we have

$$V_\alpha \cap V_\beta = \pi_\alpha^{-1}(\pi_\alpha(V_\alpha \cap V_\beta)) = \pi_\beta^{-1}(\pi_\beta(V_\alpha \cap V_\beta)).$$

(4) For each α, $\beta \in A$ satisfying $V_\alpha \cap V_\beta \neq \emptyset$ and $\alpha \neq \beta$, $U_{\alpha\beta}$ is a manifold, $\pi_{\alpha\beta} : V_\alpha \cap V_\beta \to U_{\alpha\beta}$ is a fiber bundle with fiber a closed manifold, $p^\alpha_{\alpha\beta} : U_{\alpha\beta} \to \pi_\alpha(V_\alpha \cap V_\beta)$ is a fiber bundle with fiber a closed manifold such that the following diagram commutes

$$\begin{tikzcd}
 & & V_\alpha \cap V_\beta \arrow[d, "\pi_{\alpha\beta}"] \arrow[lldd, "\pi_\alpha|_{V_\alpha\cap V_\beta}"'] \arrow[rrdd, "\pi_\beta|_{V_\alpha\cap V_\beta}"] & & \\
 & & U_{\alpha\beta} \arrow[lld, "p^\alpha_{\alpha\beta}"] \arrow[rrd, "p^\beta_{\alpha\beta}"'] & & \\
U_\alpha \supset \pi_\alpha(V_\alpha \cap V_\beta) & & & & \pi_\beta(V_\alpha \cap V_\beta) \subset U_\beta.
\end{tikzcd}$$

Here we assume $U_{\alpha\beta} = U_{\beta\alpha}$, $\pi_{\alpha\beta} = \pi_{\beta\alpha}$, and $p^\alpha_{\alpha\beta} = p^\alpha_{\beta\alpha}$.

(5) For each $b \in U_{\alpha\beta}$ we have

$$\pi_{\alpha\beta}^{-1}(b) = \pi_\alpha^{-1}(p^\alpha_{\alpha\beta}(b)) \cap \pi_\beta^{-1}(p^\beta_{\alpha\beta}(b)).$$

We often denote a compatible fibration on M by $\{\pi_\alpha\}_{\alpha \in A}$ for simplicity.

Let $\{\pi_\alpha\}_{\alpha \in A}$ be a compatible fibration on M.

Definition 2.2. For $\alpha \in A$ and $x \in V_\alpha$, we define $A(\alpha; x)$ as follows

$$A(\alpha; x) := \{\beta \in A \mid x \in V_\alpha \cap V_\beta,\ p^\beta_{\alpha\beta} : U_{\alpha\beta} \to \pi_\beta(V_\alpha \cap V_\beta) \text{ is a diffeomorphism}\}.$$

Remark 2.3. We have $\pi_\beta^{-1}\pi_\beta(x) \subset \pi_\alpha^{-1}\pi_\alpha(x)$ for $\beta \in A(\alpha, x)$.

Definition 2.4. An open covering $\{V'_\alpha\}_{\alpha \in A}$ of M is said to be *admissible* if it satisfies the following conditions

(1) $V'_\alpha \subset V_\alpha$ for each $\alpha \in A$ and

(2) $\pi_\beta^{-1}(\pi_\beta(V'_\alpha \cap V_\beta)) = V'_\alpha \cap V_\beta$ for each α and β .

Later we will fix an admissible open covering satisfying some good property. (See Assumption 2.12 (2).)

Proposition 2.5. *Let $\{V'_\alpha\}_{\alpha \in A}$ be an admissible open covering of M. Then, the collection of data*

- $\{\pi_\alpha|_{V'_\alpha} \colon V'_\alpha \to U'_\alpha := \pi_\alpha(V'_\alpha) \mid \alpha \in A\}$
- $\{\pi_{\alpha\beta}|_{V'_\alpha \cap V'_\beta} \colon V'_\alpha \cap V'_\beta \to U'_{\alpha\beta} = \pi_{\alpha\beta}(V'_\alpha \cap V'_\beta) \mid \alpha, \beta \in A$ *such that* $V'_\alpha \cap V'_\beta \neq \emptyset$ *and* $\alpha \neq \beta\}$, *and*
- $\{p^\alpha_{\alpha\beta}|_{U'_{\alpha\beta}} \colon U'_{\alpha\beta} \to \pi_\alpha(V'_\alpha \cap V'_\beta) \mid \alpha, \beta \in A$ *such that* $V'_\alpha \cap V'_\beta \neq \emptyset$ *and* $\alpha \neq \beta\}$

is a compatible fibration on M.

Suppose that there is an admissible open covering $\{V'_\alpha\}_{\alpha\in A}$ of M. We take and fix it.

Definition 2.6. A subset C of M is said to be *admissible* (for $\{V'_\alpha\}_{\alpha\in A}$) if $V'_\alpha \cap C = \pi_\alpha^{-1}(\pi_\alpha(V'_\alpha \cap C))$ for each $\alpha \in A$.

Example 2.7. Each V_α (resp. V'_α) is admissible open subset.

We can restrict a compatible fibration to an admissible open subset.

Proposition 2.8. *Let C be a submanifold of M. Suppose C is admissible. Then the data $\{\pi_\alpha|_{C\cap V'_\alpha} \colon C \cap V'_\alpha \to \pi_\alpha(C \cap V'_\alpha)\}_{\alpha\in A}$ is a compatible fibration on C.*

Proof. It is sufficient to show the following three equalities

(1) $\pi_\alpha^{-1}\pi_\alpha(C \cap V'_\alpha \cap V'_\beta) = C \cap V'_\alpha \cap V'_\beta$,
(2) $\pi_{\alpha\beta}^{-1}\pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta) = C \cap V'_\alpha \cap V'_\beta$, and
(3) ${p^\alpha_{\alpha\beta}}^{-1} p^\alpha_{\alpha\beta}(\pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta)) = \pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta)$.

First let us show the right facing inclusion $\subset$ for 1. For each $z \in \pi_\alpha^{-1}\pi_\alpha(C \cap V'_\alpha \cap V'_\beta)$ there exists $x \in C \cap V'_\alpha \cap V'_\beta$ such that $\pi_\alpha(z) = \pi_\alpha(x)$. Then $z \in \pi_\alpha^{-1}\pi_\alpha(x) \subset \pi_\alpha^{-1}\pi_\alpha(C \cap V'_\alpha) \cap \pi_\alpha^{-1}\pi_\alpha(V'_\alpha \cap V'_\beta) = C \cap V'_\alpha \cap V'_\beta$.

Next we show the right facing inclusion $\subset$ for 2. For each $z \in \pi_{\alpha\beta}^{-1}\pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta)$ there exists $x \in C \cap V'_\alpha \cap V'_\beta$ such that $\pi_{\alpha\beta}(z) = \pi_{\alpha\beta}(x)$. Then $\pi_\alpha(x) = p^\alpha_{\alpha\beta} \circ \pi_{\alpha\beta}(x) = p^\alpha_{\alpha\beta} \circ \pi_\alpha(z) = \pi_\alpha(z)$. In particular $z \in \pi_\alpha^{-1}\pi_\alpha(x) \subset \pi_\alpha^{-1}\pi_\alpha(C \cap V'_\alpha \cap V'_\beta) = C \cap V'_\alpha \cap V'_\beta$.

Finally we show the right facing inclusion $\subset$ for 3. For each $z \in {p^\alpha_{\alpha\beta}}^{-1} p^\alpha_{\alpha\beta}(\pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta))$ there exists $x \in C \cap V'_\alpha \cap V'_\beta$ such that $p^\alpha_{\alpha\beta}(z) = \pi_\alpha(x)$. We show that $\pi_{\alpha\beta}^{-1}(z) \subset C$. For $w \in \pi_{\alpha\beta}^{-1}(z)$ we have $\pi_\alpha(w) = p^\alpha_{\alpha\beta} \circ \pi_{\alpha\beta}(w) = p^\alpha_{\alpha\beta}(z) = \pi_\alpha(x)$. Then $w \in \pi_\alpha^{-1}\pi_\alpha(x) \subset C$. This shows $\pi_{\alpha\beta}^{-1}(z) \subset C$. Hence $z \in \pi_{\alpha\beta}(\pi_{\alpha\beta}^{-1}(z)) \subset \pi_{\alpha\beta}(C \cap V'_\alpha \cap V'_\beta)$. ☐

Remark 2.9. For an admissible submanifold C of V, the restriction $\{\pi_\alpha|_{C\cap V'_\alpha} \colon C \cap V'_\alpha \to \pi_\alpha(C \cap V'_\alpha)\}$ does not satisfy the second and the third conditions in Assumption 2.12 in general.

Definition 2.10. Let $f : M \to \mathbb{R}$ be a function. If f is constant along fibers of $\pi_\alpha|_{V'_\alpha}$ for all $\alpha \in A$, then we call f an *admissible function* (for $\{V'_\alpha\}_{\alpha\in A}$).

Definition 2.11. Suppose that there is an admissible open covering $\{V'_\alpha\}_\alpha$ and fix it. If a linear map $I : C^\infty(M) \to C^\infty(M)$ satisfies the following properties, then we call I an *averaging operation* for $\{V'_\alpha\}$.

(1) $I(f)$ is an admissible function for $\{V'_\alpha\}$ for all $f \in C^\infty(M)$.
(2) If f is a constant function, then $I(f)$ is also a constant function with the same value of f.
(3) If f is a non-negative function, then $I(f)$ is so.
(4) There exists an open covering $\{V''_\alpha\}_{\alpha\in A}$ of M which satisfies the following properties.
    - For all $\alpha \in A$ we have $\overline{V''_\alpha} \subset V_\alpha$.
    - For all $\alpha \in A$ we have $\pi_\alpha^{-1}\pi_\alpha(V''_\alpha) = V''_\alpha$.
    - For all $f \in C^\infty(V)$ and $x \in M$ there exists some $\alpha \in A$ such that $x \in V''_\alpha$ and

$$\min_{y\in\pi_\alpha^{-1}\pi_\alpha(x)} f(y) \le I(f)(x) \le \max_{y\in\pi_\alpha^{-1}\pi_\alpha(x)} f(y).$$

(5) Let $f : M \to \mathbb{R}$ be a smooth function. If *supp*f is contained in V'_α for some $\alpha \in A$, then *supp*$I(f)$ is also contained in V'_α.

In this paper we impose the following technical assumptions for a compatible fibration $\{\pi_\alpha : V_\alpha \to U_\alpha\}_{\alpha \in A}$.

Assumption 2.12. (1) The index set A is a finite set.

(2) There is an admissible open covering $\{V'_\alpha\}_{\alpha\in A}$ of M such that $\overline{V'_\alpha} \subset V_\alpha$, and we fix it. Hereafter we say *admissible* (resp. an *averaging operation*) instead of admissible (resp. an *averaging operation*) for $\{V'_\alpha\}$.

(3) There is an averaging operation $I : C^\infty(M) \to C^\infty(M)$ whose definition is given in Definition 2.11.

Using the averaging operation we can construct an *admissible partition of unity* as in the following.

Lemma 2.13 (Existence of admissible partition of unity)**.** *Let M be a manifold with a compatible fibration $\{\pi_\alpha\}$. There is a smooth partition of unity $\{\rho_\alpha^2\}$ of the open covering $M = \cup_\alpha V'_\alpha$ such that each ρ_α is admissible.*

Proof. Take any partition of unity $\{\phi_\alpha\}_\alpha$ of $M = \cup_\alpha V'_\alpha$. Applying the averaging operation we have a family of admissible functions $\{I(\phi_\alpha)\}_\alpha$. Note that it is an another partition of unity of $\{V'_\alpha\}$ because of the Property 2,3 and 5 of the averaging operation. We put $\rho_\alpha := I(\phi_\alpha)/\sqrt{\sum_\beta I(\phi_\beta)^2}$. Then $\{\rho_\alpha^2\}_\alpha$ is a required admissible partition of unity. □

We give a sufficient condition for Assumption 2.12.

Definition 2.14 (Good compatible fibration)**.** If a compatible fibration $\{\pi_\alpha : V_\alpha \to U_\alpha\}$ over M satisfies 1 and 2 in Assumption 2.12 together with the following 5', then we call $\{\pi_\alpha : V_\alpha \to U_\alpha\}$ a *good compatible fibration*.

5'. If $V_\alpha \cap V_\beta \neq \emptyset$, then we have $\alpha \in A(\beta; x)$ or $\beta \in A(\alpha; x)$ for all $x \in V_\alpha \cap V_\beta$.

We show the following proposition in Appendix A.

Proposition 2.15. *If $\{\pi_\alpha\}_{\alpha\in A}$ is a good compatible fibration on M, then there exists an open covering $\{V''_\alpha\}_{\alpha\in A}$ of M and a linear map $I : C^\infty(M) \to C^\infty(M)$ such that I satisfies all conditions in Definition 2.11 for $\{V''_\alpha\}_{\alpha\in A}$.*

We also give a sufficient condition for Assumption 2.12 (2) in Appendix B.

Now we define an appropriate notion of Riemannian metric and connection for a compatible fibration. For the projection π of a fiber bundle over a manifold we denote by $T[\pi]$ the vector bundle consisting of tangent vectors along fibers of π. We first note that there exist following four types of short exact sequences for $i = \alpha, \beta$

$$0 \to T[\pi_i] \to TV_i \to \pi_i^* TU_i \to 0, \tag{1}$$

$$0 \to T[\pi_{\alpha\beta}] \to T(V_\alpha \cap V_\beta) \to \pi_{\alpha\beta}^* TU_{\alpha\beta} \to 0, \tag{2}$$

$$0 \to T[\pi_{\alpha\beta}] \to T[\pi_i]|_{V_\alpha \cap V_\beta} \to \pi_{\alpha\beta}^* T[p^i_{\alpha\beta}] \to 0, \tag{3}$$

$$0 \to T[p^i_{\alpha\beta}] \to TU_{\alpha\beta} \to {p^i_{\alpha\beta}}^* T\pi_i(V_\alpha \cap V_\beta) \to 0.. \tag{4}$$

Definition 2.16. Let E^0, E^1 and E^2 be smooth vector bundles with metrics. A short exact sequence $0 \to E^0 \to E^1 \to E^2 \to 0$ is *orthogonally split* if the isomorphism $E^1 \cong$

$E^0 \oplus E^2$ defined by the orthogonal splitting with respect to the metric on E^1 is isometric with respect to the metrics on E^0, E^1 and E^2.

Definition 2.17. A *compatible Riemannian metric of a compatible fibration* is a collection of metrics on the vector bundles $T[\pi_i], TU_i$, $T[\pi_{\alpha\beta}]$, $TU_{\alpha\beta}$, and $T[p^i_{\alpha\beta}]$ such that the exact sequences (3) and (4) are orthogonally split with respect to these metrics.

From the definition, we have a canonical isometric isomorphism

$$(T[\pi_\alpha] \oplus \pi_\alpha^* TU_\alpha)|_{V_\alpha \cap V_\beta} \cong (T[\pi_{\alpha\beta}] \oplus \pi_{\alpha\beta}^* TU_{\alpha\beta}) \cong (T[\pi_\beta] \oplus \pi_\beta^* TU_\beta)|_{V_\alpha \cap V_\beta} \tag{5}$$

over $V_\alpha \cap V_\beta$.

Definition 2.18. Suppose we have a compatible fibration with a compatible Riemannian metric. A *compatible connection* is a collection of the splittings of of the short exact sequences (1) and (2) such that the isomorphism (5) is equal to the composition of the isomorphisms

$$(T[\pi_i] \oplus \pi_i^* TU_i)|_{V_\alpha \cap V_\beta} \cong T(V_\alpha \cap V_\beta) \cong (T[\pi_{\alpha\beta}] \oplus \pi_{\alpha\beta}^* TU_{\alpha\beta})$$

induced from the splittings.

2.2. **Acyclic compatible system.**

Definition 2.19. Suppose we have a compatible fibration $\{\pi_\alpha\}$ on M with a compatible Riemannian metric. A bundle W over M is a *compatible Clifford module bundle* if we have the following structures.

1. W has a structure of a $\mathbb{Z}/2$-graded $Cl(T[\pi_\alpha] \oplus \pi_\alpha^* TU_\alpha)$-module bundle over V_α.
2. Over $V_\alpha \cap V_\beta$, the above module structures on V_α and V_β are compatible with the isomorphism (5).

The next lemma follows immediately from the definitions of compatible metric, compatible connection and compatible Clifford module bundle.

Lemma 2.20. *Suppose we have a compatible metric and compatible connection. Then we have a well-defined Riemannian metric on M. Moreover if we have a compatible Clifford module bundle, then it has a structure of Clifford module with respect to the well-defined Riemannian metric on M.*

Let $\{\pi_\alpha\}$ be a compatible fibration on M with compatible Riemannian metric and $W \to M$ a compatible Clifford module bundle.

Definition 2.21 (Compatible system of Dirac-type operators)**.** A *compatible system of Dirac-type operators* on $(\{\pi_\alpha\}, W)$ is a data $\{D_\alpha\}$ satisfying the following properties.

1. $D_\alpha \colon \Gamma(W|_{V_\alpha}) \to \Gamma(W|_{V_\alpha})$ is an order-one formally self-adjoint differential operator of degree-one.
2. D_α contains only the derivatives along fibers of $\pi_\alpha \colon V_\alpha \to U_\alpha$, i.e. D_α commutes with multiplication of the pull-back of smooth functions on U_α.
3. The principal symbol $\sigma(D_\alpha)$ of D_α is given by $\sigma(D_\alpha) = c \circ p_\alpha \circ \iota_\alpha^* \colon T^*V_\alpha \to \mathrm{End}(W|_{V_\alpha})$, where $\iota_\alpha \colon T[\pi_\alpha] \to TV_\alpha$ is the natural inclusion, $p_\alpha \colon T^*[\pi_\alpha] \to T[\pi_\alpha]$ is the isomorphism induced by the Riemannian metric and $c \colon T[\pi_\alpha] \to \mathrm{End}(W|_{V_\alpha})$ is the Clifford multiplication.

(4) For $b \in U_\alpha$ and $u \in T_bU_\alpha$, let $\widetilde{u} \in \Gamma(\pi_\alpha^*TU_\alpha|_{\pi_\alpha^{-1}(b)})$ be the section naturally induced by u. $\widetilde{u}$ acts on $W|_{\pi_\alpha^{-1}(b)}$ by the Clifford multiplication $c(\widetilde{u})$. Then D_α and $c(\widetilde{u})$ anti-commute each other, i.e.

$$0 = \{D_\alpha, c(\widetilde{u})\} := D_\alpha \circ c(\widetilde{u}) + c(\widetilde{u}) \circ D_\alpha$$

for all $b \in U_\alpha$ and $u \in T_bU_\alpha$.

(5) If $V_\alpha \cap V_\beta \neq \emptyset$, then the anti-commutator $\{D_\alpha, D_\beta\} := D_\alpha \circ D_\beta + D_\beta \circ D_\alpha$ is a differential operator along fibers of $\pi_{\alpha\beta}$ of order at most 2.

The properties 1, 2, and 3 in Definition 2.21 imply that D_α is of Dirac-type when restricted to each fiber of π_α.

We call a compatible system of Dirac-type operators $\{D_\alpha\}$ a compatible system for short.

Remark 2.22. If there is a compatible system of Dirac-type operators, then, the underlying compatible fibration $\{\pi_\alpha\}_{\alpha\in A}$ must satisfies the following condition on each $V_\alpha \cap V_\beta \neq \emptyset$.

- $T[p_{\alpha\beta}^\alpha]$ and $T[p_{\alpha\beta}^\beta]$ are perpendicular to each other with respect to the metric on $TU_{\alpha\beta}$.

Definition 2.23 (Acyclic compatible system)**.** A compatible system $\{D_\alpha\}_{\alpha\in A}$ is *acyclic* if for all $\alpha \in A$, $x \in V_\alpha$ and a family of non-negative numbers $\{t_\beta\}_{\beta\in A(\alpha;x)}$ satisfying $t_\beta > 0$ for some β, the operator $\sum_{\beta\in A(\alpha;x)} t_\beta D_\beta \colon \Gamma(W|_{\pi_\alpha^{-1}(\pi_\alpha(x))}) \to \Gamma(W|_{\pi_\alpha^{-1}(\pi_\alpha(x))})$ has zero kernel. Note that the above operator is well-defined because of Remark 2.3.

Definition 2.24 (strongly acyclic compatible system)**.** A compatible system $\{D_\alpha\}$ is *strongly acyclic* if it satisfies the following conditions.

(1) For each α and $b \in U_\alpha$ $D_\alpha|_{\pi_\alpha^{-1}(b)}$ has zero kernel.

(2) If $V_\alpha \cap V_\beta \neq \emptyset$, then the anti-commutator $\{D_\alpha, D_\beta\}$ is a non-negative operator over $V_\alpha \cap V_\beta$.

We first note that the following lemma.

Lemma 2.25. *A strongly acyclic compatible system is acyclic.*

Proof. If $\{D_\alpha\}$ is strongly acyclic compatible system, then we have

$$\left(D_\alpha + \sum_{\beta\in A(\alpha;x)} \tau_\beta D_\beta\right)^2 = D_\alpha^2 + \sum \tau_\beta\{D_\alpha, D_\beta\} + \left(\sum_\beta \tau_\beta D_\beta\right)^2 \geq D_\alpha^2$$

for any family of non-negative numbers $\{\tau_\beta\}$. Suppose $\left(\sum_{\beta\in A(\alpha;x)} t_\beta D_\beta\right) s = 0$ for $s \in \Gamma(W|_{\pi_\alpha^{-1}(\pi_\alpha(x))})$. Take $\alpha_0 \in A(\alpha;x)$ so that t_{α_0} is not 0. Then we have

$$\left(D_{\alpha_0} + \sum_{\beta\in A(\alpha;x)\smallsetminus\{\alpha_0\}} (t_\beta/t_{\alpha_0})D_\beta\right) s = 0$$

and $s = 0$ by the above inequality and the first condition in Definition 2.24. □

Remark 2.26. We often need a compatible fibration $\{\pi_\alpha \colon V_\alpha \to U_\alpha\}_{\alpha\in A}$ such that U_α's are orbifolds even though M and V_α's are smooth. In fact, such a compatible fibration is necessary when we deal with a torus action. In this case, by using a uniformizing chart of U_α, we extend Definition 2.1. Instead of $\{\pi_\alpha\colon V_\alpha \to U_\alpha\}_{\alpha\in A}$ we consider the following collection of data.

- $\{q_\alpha\colon \widetilde{V}_\alpha \to V_\alpha \mid \alpha \in A\}$,
- $\{\pi_\alpha\colon \widetilde{V}_\alpha \to \widetilde{U}_\alpha \mid \alpha \in A\}$,
- $\{\mathscr{F}_\alpha \text{ on } V_\alpha \mid \alpha \in A\}$
- $\{\pi_{\alpha\beta}\colon \widetilde{V}_\alpha \times_M \widetilde{V}_\beta \to \widetilde{U}_{\alpha\beta} \mid \alpha,\beta \in A \text{ such that } V_\alpha \cap V_\beta \neq \emptyset \text{ and } \alpha \neq \beta\}$, and
- $\{p^\alpha_{\alpha\beta} : \widetilde{U}_{\alpha\beta} \to \pi_\alpha(\widetilde{V}_\alpha \times_V \widetilde{V}_\beta) \mid \alpha,\beta \in A \text{ such that } V_\alpha \cap V_\beta \neq \emptyset \text{ and } \alpha \neq \beta\}$,

where for each $\alpha, \beta \in A$, π_α, $\pi_{\alpha\beta}$, $p^\alpha_{\alpha\beta}$ are fiber bundles, q_α is a finite covering and $\mathscr{F}_\alpha$ is a foliation on V_α which satisfy appropriate compatibilities. See [5] for detail. Note that in this extension U_α's do not appear anymore. We define the above data to be *good* by the same manner as Definition 2.14. In the case where M is an orbifold, as a covering $\widetilde{V}_\alpha \to V_\alpha$ in the above extension we can adopt a uniformizing chart of V_α such that $\widetilde{V}_\alpha$ is smooth. For an orbifold version of a compatible system, consult [5].

Example 2.27. Let M be $\mathbb{R} \times S^1$ with the standard Riemannian metric and (t, θ) its standard coordinate. We introduce a compatible fibration on M by

$$\pi_\alpha\colon V_\alpha := (-\infty, 1) \times S^1 \to U_\alpha := (-\infty, 1)$$
$$\pi_\beta\colon V_\beta := (-1, \infty) \times S^1 \to U_\beta := (-1, \infty).$$

Let W be the trivial rank 2 Hermitian vector bundle $M \times \mathbb{C}^2$ on M with $\mathbb{Z}/2$-grading

$$W^0 := M \times (\mathbb{C} \times 0),\ W^1 := M \times (0 \times \mathbb{C}).$$

We define the Clifford multiplication of $Cl(TM)$ on W by

$$c(\partial_\theta) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},\ c(\partial_t) = \begin{pmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}.$$

For smooth functions $f_\alpha\colon V_\alpha \to \mathbb{R}$, $f_\beta : V_\beta \to \mathbb{R}$, let D_α, D_β be differential operators on $\Gamma(W|_{V_\alpha})$, $\Gamma(W|_{V_\beta})$ which are defined by

$$D_\alpha := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \partial_\theta + f_\alpha(t,\theta) \begin{pmatrix} 0 & \sqrt{-1} \\ -\sqrt{-1} & 0 \end{pmatrix}$$
$$D_\beta := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \partial_\theta + f_\beta(t,\theta) \begin{pmatrix} 0 & \sqrt{-1} \\ -\sqrt{-1} & 0 \end{pmatrix}.$$

They are order-one formally self-adjoint differential operators of degree-one. Then, it is easy to see that the data $\{D_\alpha, D_\beta\}$ is an acyclic compatible system if and only if f_α and f_β satisfy the following properties.

1. $f_\alpha(t,\theta) \notin \mathbb{Z}$ for any $(t,\theta) \in V_\alpha$. The same property also holds for f_β.
2. $\frac{t_\alpha f_\alpha(t,\theta) + t_\beta f_\beta(t,\theta)}{t_\alpha + t_\beta} \notin \mathbb{Z}$ for any $(t,\theta) \in V_\alpha \cap V_\beta$ and any non-negative real numbers t_α, t_β which satisfy $t_\alpha + t_\beta \neq 0$.

Example 2.28. For non-negative integers m and n satisfying $n \leq m$ let M be $\mathbb{R}^{2m-n} \times T^n$, where we regard T^n as $\mathbb{R}^n/(2\pi\mathbb{Z})^n$. Let A be an ordered set, $\{V'_\alpha\}_{\alpha\in A}$ a finite open covering of $\mathbb{R}^{2m-n}$, and $\{R_\alpha\}_{\alpha\in A}$ a family of subspaces of $\mathbb{R}^n$ spanned by rational vectors. We assume that if $\alpha < \beta$, then $R_\alpha \subset R_\beta$. We put $V_\alpha := V'_\alpha \times T^n$ and $T_\alpha := R_\alpha / R_\alpha \cap (2\pi\mathbb{Z})^n$. Define U_α to be $V'_\alpha \times T^n/T_\alpha$ and $\pi_\alpha : V_\alpha \to U_\alpha$ to be the natural projection. Then these data define a good compatible fibration on M.

Let (g, J) be the pair of the Riemannian metric and the almost complex structure on M which is defined by

$$g_x\left(\sum_{i=1}^{2m-n} a_i^1\partial_{y_i} + \sum_{i=1}^{n} b_i^1\partial_{\theta_i}, \sum_{j=1}^{2m-n} a_j^2\partial_{y_j} + \sum_{j=1}^{n} b_j^2\partial_{\theta_j}\right) = \sum_{i=1}^{2m-n} a_i^1a_i^2 + \sum_{i=1}^{n} b_i^1b_i^2,$$

$$J(\partial_{y_i}) = \begin{cases} \partial_{\theta_i} & 1 \le i \le n \\ \partial_{y_{m-n+i}} & n+1 \le i \le m \\ -\partial_{y_{i-m+n}} & m+1 \le i \le 2m-n \end{cases}$$

for $x = (y, \theta) \in M$. Note that since g is invariant under J, (g, J) defines the Hermitian metric on M by

$$h_x(u, v) = g_x(u, v) + \sqrt{-1}g_x(u, Jv)$$

for $u, v \in T_xM$. By using the horizontal lift $\pi_\alpha^*TU_\alpha \to TV_\alpha$ with respect to g, it is obvious that $\{\pi_\alpha\}$ is equipped with a compatible Riemannian metric and a compatible connection.

Next we define a compatible Clifford module bundle W and a strongly acyclic compatible system $\{D_\alpha\}_{\alpha\in A}$ in the following way. Take a Hermitian line bundle (L, ∇^L) with Hermitian connection on M whose restriction to $\pi_\alpha^{-1}(b)$ is a flat connection for each $\alpha \in A$ and $b \in U_\alpha$. We assume the following condition.

(∗) For all α and $b \in U_\alpha$ the restriction $\nabla^L|_{\pi_\alpha^{-1}(b)}$ is not trivially flat connection, i.e., its holonomy representation is non-trivial.

We define a Hermitian vector bundle W on M by

$$W := \wedge_{\mathbb{C}}^{\bullet}TM_{\mathbb{C}} \otimes L,$$

where $TM_{\mathbb{C}}$ is the tangent bundle regarded as a complex vector bundle with J. A Clifford module structure $c\colon Cl(TM) \to \mathrm{End}(W)$ is defined by

$$c(u)(\varphi) = u \wedge \varphi - u \llcorner \varphi \tag{6}$$

for $u \in TM$, $\varphi \in W$, where $\llcorner$ is the interior product with respect to h, namely,

$$v\llcorner(v_1 \wedge v_2 \wedge \cdots \wedge v_k) := \sum_{i=1}^{k}(-1)^{i-1}h(v_i, v)v_1 \wedge \hat{v}_i \wedge \cdots \wedge v_k,$$

$v_1, \ldots, v_k \in TM$.

Let $\nabla^{\wedge_{\mathbb{C}}^{\bullet}TM_{\mathbb{C}}}$ be the Hermitian connection on $\wedge_{\mathbb{C}}^{\bullet}TM_{\mathbb{C}}$ which is induced from the Levi-Civita connection on TM with respect to g. Two connections $\nabla^{\wedge_{\mathbb{C}}^{\bullet}TM_{\mathbb{C}}}$ and ∇^L define the Hermitian connection on W by

$$\nabla := \nabla^{\wedge_{\mathbb{C}}^{\bullet}TM_{\mathbb{C}}} \otimes \mathrm{id} + \mathrm{id} \otimes \nabla^L.$$

Then we define D_α by

$$\begin{aligned} D_\alpha := c \circ \iota_\alpha^* \circ \nabla|_{V_\alpha} : \Gamma\left(W|_{V_\alpha}\right) &\xrightarrow{\nabla|_{V_\alpha}} \Gamma\left(T^*V_\alpha \otimes W|_{V_\alpha}\right) \\ &\xrightarrow{\iota_\alpha^*} \Gamma\left(T^*[\pi_\alpha]) \otimes W|_{V_\alpha}\right) \\ &\xrightarrow{c} \Gamma\left(W|_{V_\alpha}\right), \end{aligned}$$

where $\iota_\alpha\colon T[\pi_\alpha] \to TV_\alpha$ is the natural inclusion.

By the construction it is obvious that $\{D_\alpha\}$ satisfies the condition 1, 2, and 3 in Definition 2.21. The condition 4 in Definition 2.21 follows from the fact that g restricted to each fiber of π_α is flat. We can show that for each α and $b \in U_\alpha$ the kernel of $D_\alpha|_{\pi_\alpha^{-1}(b)}$ vanishes. It follows from Property $(*)$ and the following lemma.

Lemma 2.29. *Let* $(E, \nabla^E) \to T$ *be a flat Hermitian line bundle on an* n*-torus. If the degree zero cohomology* $H^0(T; E)$ *with local system* (E, ∇^E) *vanishes, then all cohomologies* $H^\bullet(T; E)$ *vanish.*

Proof. Fix a flat Riemannian metric on T. Let $(x_1, \cdots, x_n)$ be a coordinate on T such that $\{\partial/\partial x_i\}$ forms a parallel orthonormal frame of the tangent bundle. Let ∇^T be the Levi-Civita connection of the flat metric. Then the induced connection $\nabla^W := \nabla^E \otimes \mathrm{id} + \mathrm{id} \otimes \nabla^T$ on $W := E \otimes \wedge^\bullet T^*T$ is a flat connection and (W, ∇^W) is a Dirac bundle in the sense of [10]. The general Bochner identity for Dirac bundles (Theorem 8.2 in [10]) implies that any harmonic section in $\Gamma(W)$ is parallel. If $\omega = \sum_I s_I dx_I \in \Gamma(W)$ is a harmonic section of degree k, where I runs multi-indices $I = (i_1 < \cdots < i_k)$ and $dx_I = dx_{i_1} \wedge \cdots \wedge dx_{i_k}$, then we have $0 = \nabla^W \omega = \sum_I (\nabla^E s_I) dx_I$, and hence, $\nabla^E s_I = 0$. Since $H^0(M; E) = 0$ we have $s_I = 0$ and $\omega = 0$. $\square$

These facts and Proposition 2.34, which will be shown in the next subsection, imply that $\{D_\alpha\}$ is a strongly acyclic compatible system.

2.3. **Example from torus action.** Suppose an n-dimensional torus G acts on a smooth compact manifold $\hat{M}$ smoothly and let M be a G-invariant open subset of $\hat{M}$. Let A be the set of all the subgroups of G which appear as stabilizers

$$G_x := \{g \in G \mid gx = x\}$$

at some points $x \in M$. Note that A is a finite set and has a partial order with respect to the inclusion.

The following lemma is useful to construct a good compatible fibration satisfying a convenient property for some cases with torus actions. We give a proof in Appendix C.

Lemma 2.30 (Existence of a good open covering)**.** *There exists an open covering* $\{V_H\}_{H \in A}$ *of* M *parameterized by* A *satisfying the following properties.*

1. *Each* V_H *is* G*-invariant.*
2. *For each* $x \in V_H$ *we have* $G_x \subset H$*.*
3. *If* $V_H \cap V_{H'} \neq \emptyset$*, then we have* $H \subset H'$ *or* $H \supset H'$*.*

Moreover if for each $x \in M$ *there is an open neighborhood* V_x *of* x *such that* $V_{gx} = gV_x$ *for all* $g \in G$*, then* $\{V_H\}_{H \in A}$ *satisfies* $V_H \subset \bigcup_{G_x = H} V_x$*.*

Proposition 2.31. *In the above case,* M *has a structure of a good compatible fibration.*

Proof. We endow G with a rational flat Riemannian metric. Precisely speaking we take a Euclidean metric on the Lie algebra of G such that the intersection of the integral lattice and the lattice generated by some orthonormal basis has rank n. We extend it to the whole G.

For a subgroup H of G let $H^\perp$ be the orthogonal complement of H defined as the image of the orthogonal complement of the Lie algebra of H by the exponential map. Since the metric is rational $H^\perp$ is well-defined as a compact subgroup of G and it has only finitely many intersection points $H \cap H^\perp$.

Let $\{V_H\}_{H \in A}$ be the open covering of M in Lemma 2.30. For each $H \in A$ we define U_H to be $V_H/H^\perp$ and $\pi_H : V_H \to U_H$ to be the natural projection. Then the data

$\{\pi_H : V_H \to U_H \mid H \in A\}$ define the good compatible fibration because of the property of the good covering. ☐

Remark 2.32. In the above proof, since the intersection $H \cap H^{\perp}$ has finitely many points the quotient space U_H may have orbifold singularies. For such a case we can construct a compatible fibration in the sense of Remark 2.26 as follows.

For each $x \in M$ let Gx be the orbit through x. Let $\{V_H\}_{H\in A}$ be the open covering of M in Lemma 2.30. For each $H \in A$ and $x \in V_H$ we take a slice $U_{H,x} \subset V_H$ at x such that $U_{H,x}$ is diffeomorphic to an open disc with dimension $\dim M - \dim Gx$ and $U_{H,x}$ intersects transversely with Gx at only one point x. Let $V_{H,x}$ be a union of all orbits which intersects with $U_{H,x}$. $V_{H,x}$ is a tubular neighborhood of Gx. We put $\widetilde{V}_{H,x} := G_x^{\perp} \times U_{H,x}$ and denote by $q_{H,x}\colon \widetilde{V}_{H,x} \to V_{H,x}$ the map which is defined by the action. $q_{H,x}$ is a finite covering. Since $x \in V_H$, by the second property of a good open covering, we have $G_x \subset H$. Then, by definition, we have $H^{\perp} \subset G_x^{\perp}$. In particular, $H^{\perp}$ acts freely on $\widetilde{V}_{H,x}$ by multiplication to the first factor. We define $\widetilde{U}_{H,x}$ to be the quotient space of the $H^{\perp}$-action on $\widetilde{V}_{H,x}$ and $\pi_{H,x}\colon \widetilde{V}_{H,x} \to \widetilde{U}_{H,x}$ to be the natural projection.

Since M is a G-invariant open subset of a compact G-manifold for each $H \in A$ we can choose finitely many points $A(H) \subset M$ such that $\{V_{H,\alpha}\}_{\alpha\in A(H)}$ covers V_H. Then, $\{q_{H,\alpha}, \pi_{H,\alpha} \mid H \in A, \alpha \in A(H)\}$ is a good compatible fibration in the sense of Remark 2.26.

Remark 2.33. We will show in Section 6 that there is an example that has a good compatible fibration, but does not have a global torus action.

2.3.1. *Family of flat torus bundles.* Let $\{\pi_\alpha\}_{\alpha\in A}$ be a compatible fibration on V with a compatible Riemannian metric and W a compatible Clifford module bundle on V. We show the following.

Proposition 2.34. *Suppose that an acyclic compatible system $\{\pi_\alpha, W, D_\alpha\}$ satisfies the following four conditions.*

- *$\pi_\alpha\colon V_\alpha \to U_\alpha$ is a good compatible fibration.*
- *$\pi_\alpha : V_\alpha \to U_\alpha$ is a flat torus bundle for all α.*
- *There is a Clifford connection ∇ on W such that the restriction of ∇ on each fiber of π_α is a flat connection for all α.*
- *D_α is the Dirac operator along fibers of π_α defined by $\nabla|_{V_\alpha}$ for all $\alpha \in A$.*

Then for all $\alpha, \beta \in A$ such that $p^{\beta}_{\alpha\beta}\colon U_{\alpha\beta} \to \pi_\beta(V_\alpha \cap V_\beta)$ is a diffeomorphism, the anti-commutator $D_\alpha \circ D_\beta + D_\beta \circ D_\alpha$ is a non-negative operator along fibers of π_β. In particular if $\ker D_\alpha = 0$ for all $\alpha \in A$, then $\{\pi_\alpha, W, D_\alpha\}$ is strongly acyclic.

It suffices to show the non-negativity of the anti-commutator along fibers. We consider the following model.

- E : Euclidean space
- Γ : maximal lattice of E
- $F := E/\Gamma$: flat torus
- $W \to F$: $Cl(TF)$-module bundle
- $\nabla : \Gamma(W) \to \Gamma(TF \otimes W)$: flat Clifford connection of W
- $c : TF \otimes W \to W$: Clifford action of TF
- A : finite set
- $\{E_\alpha\}_{\alpha\in A}$: family of subspaces of E

- $\{p_\alpha\}$: family of orthogonal projections to $\{E_\alpha\}$
- We assume $p_\alpha p_\beta = p_\beta p_\alpha$ for all $\alpha, \beta \in A$.

Using the metric we have the identification $TF = T^*F = F \times E$. For a symmetric endmorphism $S : E \to E$ let $\hat{S} : F \times E \to F \times E$ be the induced bundle map on the (co)tangent bundle. We define a differential operator D_S by the composition

$$D_S := c \circ \hat{S} \circ \nabla : \Gamma(W) \to \Gamma(W).$$

Since S is symmetric D_S is a self-adjoint operator. Let S_1 and S_2 be symmetric endmorphisms which commute each other. By the direct computation using the orthonormal basis of E consisting of simultaneously eigenvectors of S_1 and S_2 we have the following equality

$$D_{S_1} \circ D_{S_2} + D_{S_2} \circ D_{S_1} = 2\nabla^* \circ \hat{S}_1 \circ \hat{S}_2 \circ \nabla,$$

where $\nabla^* : \Gamma(TF \otimes W) \to \Gamma(W)$ is the adjoint operator of ∇.

In particular, by putting $S_1 := p_\alpha$ and $S_2 := p_\beta$ we have the following.

Lemma 2.35. *$D_\alpha D_\beta + D_\beta D_\alpha = 2D_{\alpha\beta}^2$, where $D_{\alpha\beta}$ is the self-adjoint operator $c \circ \hat{p}_{\alpha\beta} \circ \nabla$ defined by the projection $p_{\alpha\beta}$ to the intersection $E_\alpha \cap E_\beta$.*

Proof of Proposition 2.34. Since $\{\pi_\alpha\}$ is a good compatible fibration we have a family of tori at each point on V which comes from a family of subspaces whose projections commute each other. Then the claim follows from Lemma 2.35. ☐

Note that a product of good compatible fibrations is not a good compatible fibration. To formulate a product formula, we will need the following Proposition 2.36 which is a generalization of Proposition 2.34. We prepare a notation.

- Suppose $\pi_\alpha \colon V_\alpha \to U_\alpha$ is a torus bundle with structure group $GL_n(\mathbb{Z}) \ltimes \mathbb{R}^n/\mathbb{Z}^n$.
- For each $x \in V_\alpha$ and $v \in T[\pi_\alpha]_x$, we have a vector field on the fiber $\pi_\alpha^{-1}\pi_\alpha(x)$ which is parallel on the torus and is extending v.
- Using a local trivialization of the torus bundle $\pi_\alpha \colon V_\alpha \to U_\alpha$ on a neighborhood of the fiber $\pi_\alpha^{-1}\pi_\alpha(x)$, we extend the vector field to a vector field $X_\alpha(v)$ on the neighborhood.

Proposition 2.36. *Suppose that an acyclic compatible system $\{\pi_\alpha, W, D_\alpha\}$ satisfies the following three conditions.*

- *For $x \in V_\alpha \cap V_\beta$, $v_\alpha \in T[\pi_\alpha]_x$ and $v_\beta \in T[\pi_\beta]_x$, we have $[X_\alpha(v_\alpha), X_\beta(v_\beta)] = 0$. Moreover, if $v_\alpha = v_\beta$, then, $X_\alpha(v_\alpha)$ and $X_\beta(v_\beta)$ coincide on a neighborhood of $x \in M$.*
- *There is a Clifford connection ∇ on W such that the restriction of ∇ on each fiber of π_α is a flat connection for all α.*
- *D_α is the Dirac operator along fibers of π_α defined by $\nabla|_{V_\alpha}$ for all $\alpha \in A$.*

Then for all $\alpha, \beta \in A$ such that $p^\beta_{\alpha\beta} \colon U_{\alpha\beta} \to \pi_\beta(V_\alpha \cap V_\beta)$ is a diffeomorphism, the anti-commutator $D_\alpha \circ D_\beta + D_\beta \circ D_\alpha$ is a non-negative operator along fibers of π_β. In particular if $\ker D_\alpha = 0$ *for all $\alpha \in A$, then $\{\pi_\alpha, W, D_\alpha\}$ is strongly acyclic.*

Proof. We can use the same local model used in the proof of Proposition 2.34. ☐

2.3.2. *Symplectic manifold with a torus action.* Let (M, ω) be a $2m$-dimensional symplectic manifold equipped with a Hamiltonian action of an n-dimensional torus G. In this case each orbit is an affine isotropic torus in M. Suppose that there is a G-equivariant prequantizing line bundle (L, ∇) on (M, ω), i.e., L is a Hermitian line bundle over M with a Hermitian connection ∇ whose curvature form is equal to $-2\pi\sqrt{-1}\omega$, and all these data are G-equivariant. Since an orbit is isotropic the restriction of (L, ∇) on each orbit is a flat line bundle. Let $A = \{H\}$ be the set of all subgroups of G which appear as a stabilizer subgroup of G-action at some point in M. We ssume that A is a finite set. We show the following.

Proposition 2.37. *Suppose that for each $x \in M$ there exists an open set V_x such that for all $y \in V_x$ the restriction of L on $G_x^{\perp}$-orbit $G_x^{\perp} \cdot y$ has no nontrivial parallel sections. Then M is equipped with a strongly acyclic compatible system parametrized by A.*

Remark 2.38. The assumption of Proposition 2.37 implies that there is no global non-zero parallel section on each G-orbit. However the converse does not hold in general. On the other hand in the case of Hamiltonian torus action, the moment map image of a G-orbit with a global non-zero parallel section on it is a lattice point in the dual of the Lie algebra. In this case it may be expected that there would exist a strongly acyclic compatible system on the complement of the inverse image of lattice points. If it were the case, we would have a localization of the Riemann-Roch number as a sum of contributions from lattice points. Proposition 2.37, however, does not realize this expectation. Actually the complement of the inverse image of the lattice points does not satisfy the assumption of Proposition 2.37 in general. We would need to delete neighborhoods of some extra points from the complelent to make it satisfy the assumption. In fact it is not hard to check that the contributions from the extra points to the Riemann-Roch number turn out to be zero, for instance, in the case of toric action, which implies that the localization to the lattice points is practically realized.

Proof of Proposition 2.37. By taking the union along G-orbits we may assume that each V_x is G-invariant. According to Lemma 2.30, and Remark 2.31, we have a good compatible fibration on M using a good open covering $\{V_H\}$ parameterized by $A = \{H\}$. Note that each V_H is contained in $\cup_{G_x = H} V_x$. Fix a G-invariant ω-compatible almost complex structure J on M. Then using the associated G-invariant metric g_J we can construct a compatible Riemannian metric on M as follows. It is sufficient to construct metrics of $T[\pi_H]$ for each $H \in A$ and $T[p_{HK}]$ for a sequence $H \subset K$, and check the compatibility coming from (2) and (3). The metric on $T[\pi_H]$ is defined as the restriction of g_J, and the metric on $TU_H = T(V_H/H^{\perp})$ is defined as the quotient metric of g_J. Note that the fiber of p_{HK} is the quotient of the $H^{\perp}$-orbit by the $K^{\perp}$-orbit. Then the metric on $T[p_{HK}]$ is defined as the quotient of the restriction of the metric g_J on $H^{\perp}$-orbit by the $K^{\perp}$-action. It is straightforward to check the compatibilities. We remark that the metric g_J restricted to each orbit is a flat affine metric because it is G-invariant.

Let W be the $\mathbb{Z}/2$-graded compatible Clifford module bundle

$$W = \wedge^{\bullet}_{\mathbb{C}} TM_{\mathbb{C}} \otimes L$$

with Clifford module structure $c \colon Cl(TM) \to \mathrm{End}(W)$ defined by (6). We show that W is a compatible Clifford module bundle with some more additional structures. For $H \in A$, let $(T[\pi_H] \oplus JT[\pi_H])^{\perp}$ be the orthogonal complement of $T[\pi_H] \oplus JT[\pi_H]$ with respect to g_J. Since (g_J, J) is G-invariant $H^\perp$-action preserves $(T[\pi_H]\oplus JT[\pi_H])^\perp$ and $JT[\pi_H]$. Then we define

$$\begin{aligned}E_H &:= (T[\pi_H]\oplus JT[\pi_H])^\perp/H^\perp,\\ F_H &:= JT[\pi_H]/H^\perp.\end{aligned}$$

It is obvious that TU_H has the natural isometry $TU_H\cong E_H\oplus F_H$.

Since g_J is invariant under J, J preserves $(T[\pi_H]\oplus JT[\pi_H])^\perp$. In particular $T[\pi_H]\oplus JT[\pi_H]$ and $(T[\pi_H]\oplus JT[\pi_H])^\perp$ have the structures of Hermitian vector bundles with respect to the restriction of (g_J, J) to them. Moreover (g_J, J) is G-invariant it descends to the Hermitian structure on E_H. Then, we define $W_{1,H}$ and $W_{2,H}$ by

$$\begin{aligned}W_{1,H} &:= \wedge^\bullet_{\mathbb{C}}(T[\pi_H]\oplus JT[\pi_H])_{\mathbb{C}}\otimes L,\\ W_{2,H} &:= \wedge^\bullet_{\mathbb{C}}(E_H)_{\mathbb{C}}\otimes L,\end{aligned}$$

and define the Clifford module structures

$$\begin{aligned}c_{1,H}&\colon Cl(T[\pi_H]\oplus \pi_H^*F_H)\to \mathrm{End}\,(W_{1,H})\,,\\ c_{2,H}&\colon Cl\,(E_H)\to \mathrm{End}\,(W_{2,H})\end{aligned}$$

by the same formula as in (6). By definition, TV_H has a decomposition

$$TV_H=(T[\pi_H]\oplus\pi_H^*F_H)\oplus\pi_H^*E_H$$

as Hermitian vector bundles. With respect to this decomposition there are the following isomorphisms

$$\begin{aligned}Cl(TM)|_{V_H} &\cong Cl(T[\pi_H]\oplus\pi_H^*F_H)\otimes Cl(\pi_H^*E_H),\\ &\cong Cl(T[\pi_H]\oplus\pi_H^*TU_H)\\ W|_{V_H} &\cong W_{1,H}\otimes\pi_H^*W_{2,H}.\end{aligned}$$

Then by the direct calculation one can check $c=c_{1,H}\otimes c_{2,H}$ under the above identifications.

Now we define a strongly acyclic compatible system $\{D_H\}$ on W. Let $\nabla^{T[\pi_H]}\colon \Gamma(TV_H)\to \Gamma(T^*[\pi_H]\otimes TV_H)$ be the the family of Levi-Civita connections along fibers of π_H, namely,

$$\nabla^{T[\pi_H]}=\iota_H^*\otimes q_H\circ\nabla^{TM}\circ q_H,$$

where $\iota_H\colon T[\pi_H]\to TV_H$ is the natural inclusion, ∇^{TM} is the Levi-Civita connection on TM with respect to g_J, and $q_H\colon TV_H\to TV_H$ is the orthogonal projection to $T[\pi_H]$ with respect to g_J. $\nabla^{T[\pi_H]}$ induces the family of Hermitian connections on $\wedge^\bullet_{\mathbb{C}}TM_{\mathbb{C}}|_{V_H}$ along fibers of π_H, which is denoted by $\nabla^{\wedge^\bullet_{\mathbb{C}}TM_{\mathbb{C}}|_{V_H}}$. We define the family of Hermitian connections ∇^H on $W|_{V_H}$ along fibers of π_H by

$$\nabla^H:=\nabla^{\wedge^\bullet_{\mathbb{C}}TM_{\mathbb{C}}|_{V_H}}\otimes\mathrm{id}+\mathrm{id}\otimes\left(\iota_H^*\otimes\mathrm{id}\circ\nabla^L\right)\colon\Gamma(W|_{V_H})\to\Gamma(T^*[\pi_H]\otimes W|_{V_H}).$$

Then we define $D_H\colon\Gamma(W|_{V_H})\to\Gamma(W|_{V_H})$ to be the family of de Rham operators along fibers of π_H which is defined by

$$D_H:=c_{1,H}\circ p_H\circ\nabla^H,$$

where $p_H\colon T^*[\pi_H]\to T[\pi_H]$ is the isomorphism via g_J.

The restriction of L on each $H^\perp$-orbit has no non-trivial parallel sections because of the property $V_H\subset\cup_{G_x=H}V_x$ and our assumption. It implies $\ker D_H=0$ by Lemma 2.29. Moreover since the collection of data $\{\pi_H: V_H\to U_H, W, D_H\}$ satisfies the assumptions in Proposition 2.34, $\{D_H\}$ is strongly acyclic. □

Remark 2.39. In the above proof we used the isomorphism between $\mathbb{Z}/2$-graded Clifford algebras

$$Cl(T_1 \oplus T_2) \cong Cl(T_1) \otimes Cl(T_2)$$

for two real vector bundles T_1 and T_2 with inner products. Our convention is as follows. Let $A = A^0 \oplus A^1$ and $B = B^0 \oplus B^1$ be two $\mathbb{Z}/2$-graded algebras. Then we define a structure of a $\mathbb{Z}/2$-graded algebra on the tensor product $A \otimes B$ as follows. The $\mathbb{Z}/2$-grading is defined as the one for vector spaces,

$$A \otimes B = ((A^0 \otimes B^0) \oplus (A^1 \otimes B^1)) \oplus ((A^0 \otimes B^1) \oplus (A^1 \otimes B^0)).$$

The multiplicative structure is defined by

$$(a \otimes b) \cdot (a' \otimes b') = (-1)^{\deg b \deg a'}(aa') \otimes (bb'),$$

where $a, a' \in A^0 \cup A^1$ and $b, b' \in B^0 \cup B^1$. Now let R_A and R_B be $\mathbb{Z}/2$-graded A and B modules respectively. Then we define a structure of a $\mathbb{Z}/2$-graded $A \otimes B$-module on the tensor product $R_A \otimes R_B$ by the following formula.

$$(a \otimes b) \cdot (r_A \otimes r_B) = (-1)^{\deg b \deg r_A}(ar_A \otimes br_B),$$

where $a \in A^0 \cup A^1$, $b \in B^0 \cup B^1$, $r_A \in R_A^0 \cup R_A^1$ and $r_B \in R_B^0 \cup R_B^1$.

3. An index theory for complete Riemannian manifolds

3.1. Formulation of index on complete manifolds.

Suppose M is a complete Riemannian manifold, W is a $\mathbb{Z}/2$-graded Hermitian vector bundle, and $\sigma : TM \to \mathrm{End}(W)$ is a homomorphism such that $\sigma(v)$ is a skew-Hermitian isomorphism of degree-one for each $v \in TM \smallsetminus \{0\}$. Let D be a degree-one formally self-adjoint first-order elliptic differential operator on W with principal symbol σ. We assume that σ and the coefficients of D are smooth. We formulate an index theory on M under the following assumption.

Assumption 3.1.
- D has finite propagation speed: there exists a positive real number C_0 satisfying $|\sigma| \leq C_0$ uniformly on M,
- There exist a positive real number $\lambda_0 > 0$ and an open subset V of M with its complement $M \smallsetminus V$ compact such that

$$\lambda_0 ||s||_V^2 \leq ||Ds||_V^2$$

for any smooth compactly-supported section s of W with support contained in V.

It is known that the finite propagation speed implies that D is essentially self-adjoint [2]. We will give a direct proof of the following theorem.

Theorem 3.2. *D is essentially self-adjoint as an operator on L^2-sections of W and its spectrum is discrete in $(-\sqrt{\lambda_0}, \sqrt{\lambda_0})$.* □

The proof of the first part is given as Lemma 3.11 in Section 3.2. The rest of the statement follows from Proposition 3.14 in Section 3.3.

Definition 3.3. E_λ is the vector space of smooth sections s of W such that s is L^2-bounded and satisfies $D^2 s = \lambda s$.

Theorem 3.2 implies that E_λ is zero for $\lambda < 0$, and E_λ is finite dimensional for $\lambda < \lambda_0$. Moreover there are only discrete values $\lambda < \lambda_0$ for which E_λ is non-zero. Note that the super dimension of E_λ is zero for $0 < \lambda < \lambda_0$, and hence the super dimension of $\oplus_{\lambda < \lambda_1} E_\lambda$ is constant for $0 < \lambda_1 < \lambda_0$.

Definition 3.4. $\operatorname{ind} D$ is the super dimension of E_0, or the super dimension of $\oplus_{\lambda<\lambda_1} E_\lambda$ for $0 < \lambda_1 < \lambda_0$.

The index has the following deformation invariance. Let $\{D_t\}$ ($|t| < \epsilon$) be a one-parameter family of degree-one formally self-adjoint first-order elliptic differential operators on W with principal symbols $\{\sigma_t\}$.

Assumption 3.5.

- Each D_t and σ_t satisfy Assumption 3.1 for common λ_0 and V.
- On each compact subset of M the coefficients of D_t are C^∞ convergent to those of D_0 as $t \to 0$.

We do not assume that the propagation speed is uniform with respect to t.

We will show the following theorem in Section 3.4.

Theorem 3.6. *Under Assumption 3.5* $\operatorname{ind} D_t$ *is constant with respect to* t.

Remark 3.7. So far we are fixing M and W. In Section 3.5 we will formulate a deformation for which M and W can vary. We give a proof of Theorem 3.6 so that it can be directly generalized to this case. The generalization immediately implies an excision property of index for complete Riemannian manifolds.

3.2. **Partial integration.**

We need two partial integration formulas. In general let W be a Hermitian vector bundle over a complete Riemannian manifold M, and $D_\tau : \Gamma(W) \to \Gamma(W)$ be a first order partial differential operator on W with smooth coefficients whose principal symbol is τ. We assume that D_τ has finite propagation speed, i.e., τ is a smooth L^∞-bounded section of $TM \otimes \operatorname{End}(W)$.

Lemma 3.8. *Let* $s \in \Gamma(W)$ *is an* L^2*-bounded section such that* $D_\tau^* D_\tau s$ *is also* L^2*-bounded. Then* $D_\tau s$ *is also* L^2*-bounded and we have*

$$\int_M (D_\tau^* D_\tau s, s) = \int_M |D_\tau s|^2.$$

Lemma 3.9. *Suppose* s_0 *and* s_1 *are* L^2*-bounded sections of* W *such that* $D_\tau s_0$ *and* $D_\tau^* s_1$ *are also* L^2*-bounded. Then we have*

$$\int_M (D_\tau s_0, s_1) = \int_M (s_0, D_\tau^* s_1).$$

We follow the argument in [8] using a family of cut-off functions:

Lemma 3.10. *Let* M *be a complete Riemmanian manifold.*

1. *There is a smooth proper function* $f : M \to \mathbb{R}$ *such that* $|df|$ *is bounded and* $f^{-1}((-\infty, c])$ *is compact for any* c.
2. *There is a constant* $C > 0$ *such that for each* $\epsilon > 0$ *and* $a \in \mathbb{R}$, *we have a compact supported function* $\rho_{a,\epsilon} : M \to [0,1]$ *which is equal to* 1 *on* $f^{-1}((-\infty, a])$, *and satisfies* $|d\rho_{a,\epsilon}| < C\epsilon$.

A proof of the above lemma is given in [7]. For completeness we give a detailed construction in Appendix D. The existence of such a function in 1 of Lemma 3.10 is equivalent to the completeness of M. For mored details see [7].

If we choose a Hermitian connection ∇ on W, we can describe $D_\tau s = \tau \cdot \nabla s + Bs$, where B is a smooth section of $\operatorname{End}(W)$ and $\cdot$ is the combination of the product $\operatorname{End}(W) \otimes W \to W$ and the contraction $T^*M \otimes TM \to \underline{\mathbb{R}}$. We do not assume that B is bounded.

Proof of Lemma 3.8. We first assume that s is smooth. We follow Gromov's proof of [8, Lemma 1.1 B]. From the equality

$$\begin{aligned}\int_M (D_\tau^* D_\tau s,\ \rho_{a,\epsilon}^2 s) &= \int_M (D_\tau s,\ D_\tau(\rho_{a,\epsilon}^2 s)) \\ &= \int_M (D_\tau s,\ \rho_{a,\epsilon}^2 D_\tau s) + \int_M (D_\tau s,\ 2\rho_{a,\epsilon}\tau(d\rho_{a,\epsilon})s),\end{aligned} \tag{7}$$

there is a constant C independent of s, a, ϵ such that

$$||D_\tau^* D_\tau s||_2 ||s||_2 \geq ||\rho_{a,\epsilon} D_\tau s||_2^2 - C\epsilon ||\rho_{a,\epsilon} D_\tau s||_2.$$

It implies that, as a increases, $||\rho_{a,\epsilon} D_\tau s||_2$ is bounded, i.e., $D_\tau s$ is L^2-bounded. Using (7) again we have

$$\int_M (D_\tau^* D_\tau s, s) = ||D_\tau s||_2^2 + I, \qquad |I| \leq C\epsilon ||D_\tau s||_2 ||s||_2.$$

Taking $\epsilon \to 0$, we obtain the required equality.

When s is not smooth, take a smooth compactly supported section which approximate s in L_2^2-norm on the support of $\rho_{a,\epsilon}$. Then we can reduce the argument to the smooth case.

□

Proof of Lemma 3.9. We first assume that s is smooth. We have

$$0 = \int_M (D_\tau(\rho_{a,\epsilon} s_0), s_1) - \int_M (s_0, D_\tau^*(\rho_{a,\epsilon} s_1)) = \int_M (D_\tau s_0, s_1) - \int_M (s_0, D_\tau^* s_1) + I'$$

with an error term I' satisfying $|I'| \leq C\epsilon ||s_0||_2 ||s_1||_2$, which implies the required equality. When s is not smooth, we can reduce the argument to the smooth case as in the proof of Lemma 3.8. □

Using the cut off function and a standard argument we can also show:

Lemma 3.11. *Under Assumption 3.1 D is essentially self-adjoint.*

Proof of Lemma 3.11. Suppose L^2-sections u and v satisfies $Du = v$ weakly. We show that for any $\epsilon > 0$ there is a compactly supported smooth section u_ϵ satisfying $||u_\epsilon - u||_M \leq 2\epsilon$ and $||Du_\epsilon - v||_M \leq 4\epsilon$. Take a compact subset K such that $||u||_{M\smallsetminus K}, ||v||_{M\smallsetminus K} < \epsilon$. Using Lemma 3.10 choose a smooth compactly-supported function $\rho : M \to [0,1]$ satisfying $\rho = 1$ on K and $|d\rho| < 1$. Let K' be the compact support of ρ. Since D is elliptic, the weak equality $Du = v$ and the regularity theorem imply that u is locally L_1^2-bounded and there is a smooth sections u' satisfying $||u' - u||_{K'} < \epsilon$ and $||Du' - v||_{K'} < \epsilon$. Then we have

$$||\rho u' - u||_M \leq ||\rho(u' - u)||_{K'} + ||(1-\rho)u||_{M\smallsetminus K} \leq 2\epsilon$$

and

$$\begin{aligned}||D(\rho u') - v||_M &= ||d\rho \cdot (u' - u) + d\rho \cdot u + \rho(Du - v) - (1-\rho)v||_M \\ &\leq ||u' - u||_{K'\smallsetminus K} + ||u||_{K'\smallsetminus K} + ||Du' - v||_{K'} + ||v||_{M\smallsetminus K} \\ &\leq 4\epsilon.\end{aligned}$$

It implies that D is essentially self-adjoint. □

3.3. **Min-max principle.** In this section we use Assumption 3.1 for a single operator D, and Assumption 3.5 for a one-parameter family $\{D_t\}$.

Lemma 3.12. *For any compact subset K containing $M \smallsetminus V$ there is a compact set K' containing K such that if s and Ds are L^2-bounded, then we have the estimate*

$$\lambda_0^{1/2}||s||_{M\smallsetminus K} - 2\lambda_0^{1/2}||s||_{K'\smallsetminus K} \leq ||Ds||_{M\smallsetminus K}.$$

Moreover if the coefficients of D_t are C^∞-convergent to those of $D_0 = D$ on any compact set as $t \to 0$, then we can choose K' so that the above estimate is valid for any t sufficiently close to 0.

Proof. Lemma 3.11 implies that we can assume that s is smooth and compactly supported without loss of generality. From Lemma 3.10, for any $\epsilon > 0$, there is a compact set K' containing K and a smooth non-negative function $\rho : M \to \mathbb{R}$ such that $\rho = 1$ on $M \smallsetminus K'$, $\rho = 0$ on K and $|d\rho| \leq \epsilon$. Then the above estimate follows from the next two inequalities

$$\begin{aligned} ||D(\rho s)||_M &\geq \lambda_0^{1/2}||\rho s||_V \geq \lambda_0^{1/2}||s||_{M\smallsetminus K'} \geq \lambda_0^{1/2}||s||_{M\smallsetminus K} - \lambda_0^{1/2}||s||_{K'\smallsetminus K}, \\ ||D(\rho s)||_M &\leq ||\rho Ds||_M + ||\sigma \cdot ((d\rho)\otimes s)||_M \leq ||Ds||_{M\smallsetminus K} + C(D,K')\epsilon||s||_{K'\smallsetminus K}, \end{aligned}$$

where $C(D,K') := \max_{K'}|\sigma|$. The last statement of the lemma follows from the fact that $C(D_t, K')$ is continuous with respect to t. □

Proposition 3.13. *Suppose $0 \leq \lambda_1 < \lambda_0$. Let $\{s_i\}$ be a sequence of L^2-sections of W satisfying $||s_i||_M = 1$, and $\{t_i\}$ is a sequence convergent to 0. Suppose each $D_{t_i}s_i$ is L^2-bounded and satisfies $||D_{t_i}s_i||_M^2 \leq \lambda_1$. Then there is a subsequence $\{s_{i'}\}$ which is weakly convergent to some non-zero $s_\infty \neq 0$ such that $D_0 s_\infty$ is L^2-bounded and satisfies*

$$||D_0 s_\infty||_M^2 \leq \lambda_1 ||s_\infty||_M^2 \tag{8}$$

Proof. Take a subsequence $\{s_{i'}\}$ so that $\{s_{i'}\}$ and $\{D_{t_{i'}}s_{i'}\}$ are weakly convergent to some s_∞ and u_∞ in $L^2(M,W)$ respectively. Since D_t is a smooth family, for each smooth compactly-supported section ϕ the sequence $D_{t_{i'}}\phi$ is strongly convergent to $D_0\phi$. The equality $\int_M (D_{t_{i'}}\phi, s_{i'}) = \int_M(\phi, D_{t_{i'}}s_{i'})$ implies $\int_M(D_0\phi, s_\infty) = \int_M(\phi, u_\infty)$, i.e., $D_0 s_\infty = u_\infty$ weakly.

Since $\{D_{t_{i'}}s_{i'}\}$ is L^2-bounded, Assumption 3.5 and a priori estimate imply that on any compact set $s_{i'}$ is strongly L^2-convergent to s_∞.

On the other hand for any compact set K containing $M \smallsetminus V$ there exists a compact set K' such that

$$\lambda_0^{1/2}||s_{i'}||_{M\smallsetminus K} - 2\lambda_0^{1/2}||s_{i'}||_{K'\smallsetminus K} \leq ||D_{t_{i'}}s_{i'}||_{M\smallsetminus K} \leq \lambda_1^{1/2}$$

by Lemma 3.12. If s_∞ is 0, then we have $||s_{i'}||_{K'\smallsetminus K}$ converges to 0, which contradicts to $||s_{i'}||_M = 1$ and $\lambda_1 < \lambda_0$.

Suppose the estimate (8) does not hold. Then for any $\epsilon > 0$ and any sufficiently small $\epsilon' > 0$ there exists a compact set K containing $M \smallsetminus V$ satisfying $||s_\infty||_{M\smallsetminus K} < \epsilon$ and $\lambda_1||s_\infty||_K^2 + \epsilon' < ||D_0 s_\infty||_K^2$. We choose ϵ and ϵ' so that they satisfy $8\epsilon\lambda_0(1+2\epsilon) < \epsilon'/2$. Note that the weak convergence implies $||D_0 s_\infty||_K^2 \leq \liminf_{i'\to\infty} ||D_{t_{i'}}s_{i'}||_K^2$. Since $s_{i'}$ is strongly L^2-convergent to s_∞ on the compact set K, we have

$$\lambda_1||s_{i'}||_K^2 + \frac{\epsilon'}{2} < ||D_{i'}s_{i'}||_K^2 \tag{9}$$

for sufficiently large i'. Let K' be the compact set containing K which gives the estimate in Lemma 3.12 for sufficiently small t. Since , $s_{i'}$ is strongly L^2-convergent to s_∞ on the pre-compact set $K' \smallsetminus K$, we have $||s_{i'}||_{K' \smallsetminus K} < 2\epsilon$ for sufficiently large i'. The estimate in Lemma 3.12 implies that we have $\lambda_0^{1/2}||s_{i'}||_{M\smallsetminus K} \leq ||Ds_{i'}||_{M\smallsetminus K} + 4\epsilon\lambda_0^{1/2}$ for sufficiently large i'. Taking square, and using $\lambda_1 < \lambda_0$ and $||D_{i'}s_{i'}||_{M\smallsetminus K} \leq \lambda_1^{1/2}$, we obtain

$$\lambda_1||s_{i'}||^2_{M\smallsetminus K} \leq ||D_{i'}s_{i'}||^2_{M\smallsetminus K} + 8\epsilon\lambda_0(1+2\epsilon)$$

Adding with (9) we have $\lambda_1||s_{i'}||^2_M < ||D_{i'}s_{i'}||^2_M$ for sufficiently large i', which contradicts our assumption. ☐

For a single operator D we have

Proposition 3.14. (1) *Suppose $\lambda < \lambda' < \lambda_0$.*
   (a) *If $s \in E_\lambda$, then Ds is L^2-bounded and $||Ds||^2_M = \lambda||s||^2_M$.*
   (b) *E_λ and $E_{\lambda'}$ are L^2-orthogonal to each other.*

(2) *Suppose $0 \leq \lambda_1 < \lambda_0$.*
   (a) dim $\oplus_{\lambda\leq\lambda_1}E_\lambda(D) < \infty$.
   (b) *Let R_{λ_1} be the set of L^2-bounded sections s satisfying $||s||_M = 1$ and $||Ds||^2_M < \lambda_0$ such that s is L^2-orthogonal to $\oplus_{\lambda\leq\lambda_1}E_\lambda(D)$. If R_{λ_1} is not empty, then the functional $I_{\lambda_1} : R_{\lambda_1} \to [0, \lambda_0)$, $I_{\lambda_1}(s) = ||Ds||^2_M$ attains its minimum value.*
   (c) *Let $\lambda_2 = ||Ds_0||^2_M$ be the minimum value of I_{λ_1} at a minimum s_0. Then we have $\lambda_1 < \lambda_2 < \lambda_0$ and $s_0 \in E_{\lambda_2}$*

Proof. The first statement for $\lambda < \lambda' < \lambda_0$ follows from the partial integration formulas Lemma 3.8 and Lemma 3.9.

Suppose $\lambda_1 < \lambda_0$. If $\oplus_{\lambda\leq\lambda_1}E_\lambda(D)$ is not finite dimensional, then we have a sequence e_i in the infinite space with $||e_i||_M = 1$ and mutually L^2-orthogonal each other. Proposition 3.13 implies that we have a weakly convergent limit for a subsequence with non zero limit, which is a contradiction.

Suppose s_i is a sequence in R_{λ_1} such that $I_{\lambda_1}(s_i)$ convergent to the infimum of I_{λ_1}. Proposition 3.13 implies that we have a weakly convergent limit $s_\infty \neq 0$ for a subsequence such that $s_0 := s_\infty/||s_\infty||$ is an element of R_λ which attains the infimum. For any compactly-supported smooth section s', let s'' be the L^2-orthogonal projection of s' to $\oplus_{\lambda\leq\lambda_1}E_\lambda(D)$ and put $s''' = s' - s''$. Since s_0 attains the minimum of I_{λ_1}, the derivative of $(s_0 + ts''')/||s_0 + ts'''||_M$ at 0 vanishes, and we obtain

$$\int_M (Ds_0, Ds''') = \lambda_2 \int_M (s_0, s''').$$

Since $Ds''' = Ds' - Ds''$ and $D^2s''' = D^2s' - D^2s''$ is L^2-bounded, Lemma 3.9 implies

$$\int_M (s_0, D^2s''') = \lambda_2 \int_M (s_0, s''').$$

On the other hand we have

$$\int_M (s_0, D^2s'') = \int_M (s_0, s'') = 0.$$

These relations imply

$$\int_M (s_0, D^2s') = \lambda_2 \int_M (s_0, s'),$$

i.e., $D^2 s_0 = \lambda_2 s_0$ weakly. Lemma 3.8 implies that Ds_0 is L^2-bounded and $||Ds_0||_M^2 = \lambda_2 ||s_0||_M = \lambda_2$. Since s_0 is L^2-orthogonal to $\oplus_{\lambda \le \lambda_1} E_\lambda$, we have $\lambda_1 < \lambda_2$. The regularity theorem implies s_0 is smooth and hence $s_0 \in E_{\lambda_2}$. ☐

Corollary 3.15. *Suppose $\lambda_1 < \lambda_0$. Let E be a $\mathbb{Z}/2$ graded subspace of $L^2(M, W)$ such that such that Ds is in $L^2(M, W)$ and $||Ds||_M^2 \le \lambda_1 ||s||_M^2$ for any $s \in E$. Then E is finite dimensional and*

$$\dim \oplus_{\lambda \le \lambda_1} E_\lambda(D) \ge \dim E.$$

Moreover the above inequality holds for each degree of $\mathbb{Z}/2$.

3.4. **Deformation invariance of index.** For a family $\{D_t\}$ we have:

Proposition 3.16. *Suppose $\lambda_1 < \lambda_0$. Let $\{t_i\}$ be a sequence convergent to 0 and $E^{(i)}$ be a $\mathbb{Z}/2$ graded subspace of $L^2(M, W)$ such that $D_{t_i} s$ is in $L^2(M, W)$ and $||D_{t_i} s||_M^2 \le \lambda_1 ||s||_M^2$ for any i and $s \in E_i$. Then each $E^{(i)}$ is finite dimensional and*

$$\dim \oplus_{\lambda \le \lambda_1} E_\lambda(D_0) \ge \limsup_{i \to \infty} \dim E^{(i)}.$$

Moreover the above inequality holds for each degree of $\mathbb{Z}/2$.

Proof. Suppose $\dim \oplus_{\lambda \le \lambda_1} E_\lambda(D_0) < \dim E^{(i')}$ for a subsequence $\{i'\}$. Let $s_{i'}$ be an element of $E^{(i')}$ with $||s_{i'}||_M = 1$ which is L^2-orthogonal to $\oplus_{\lambda \le \lambda_1} E_\lambda(D_0)$. Let s_∞ be the L^2-bounded section given by Proposition 3.13. Then the weak limit s_∞ is also L^2 orthogonal to $\oplus_{\lambda \le \lambda_1} E_\lambda(D_0)$, which contradicts Proposition 3.14. ☐

Remark 3.17. In the above proof the choice of $s_{i'}$ can be generalized as follows: Fix an L^2-orthonormal basis $e_1, e_2, \dots, e_N$ of the finite dimensional space $\oplus_{\lambda \le \lambda_1} E_\lambda(D_0)$. Fix any sequence $\{e_k^{(i')}\}_{i'}$ for each $1 \le k \le N$ which is strongly L^2-convergent to e_k as $i' \to \infty$. Let $s_{i'}$ be an element of $E^{(i')}$ with $||s_{i'}||_M = 1$ which is L^2-orthogonal to all $e_k^{(i')}$ $(1 \le k \le N)$. Then the rest of the proof remains valid.

Corollary 3.18. *For each degree of $\mathbb{Z}/2$ we have the inequality*

$$\dim \oplus_{\lambda \le \lambda_1} E_\lambda(D_0) \ge \limsup_{t \to 0} \dim \oplus_{\lambda \le \lambda_1} E_\lambda(D_t)$$

Proposition 3.19. *Suppose $\lambda_1 < \lambda_0$. For each degree of $\mathbb{Z}/2$ we have the inequality*

$$\dim \oplus_{\lambda < \lambda_1} E_\lambda(D_0) \le \liminf_{t \to 0} \dim \oplus_{\lambda < \lambda_1} E_\lambda(D_t)$$

Proof. Let $\epsilon_0 > 0$ be a sufficiently small number, which we fix later. Let $e_1, e_2, \dots, e_N$ be an L^2-orthonormal basis of the finite dimensional space $E := \oplus_{\lambda < \lambda_1} E_\lambda(D_0)$ consisting of eigenvectors of D_0 with eigenvalues $\mu_1, \mu_2, \dots, \mu_N$ respectively. For a sufficiently large a and sufficiently small $\epsilon > 0$, the truncated sections $e_i' = \rho_{a,\epsilon} e_i$ satisfy

$$\left| \delta_{ij} - \int_M (e_i', e_j') - \right| < \epsilon_0, \quad \left| \mu_i \mu_j \delta_{ij} - \int_M (D_0 e_i', D_0 e_j') \right| < \epsilon_0$$

for every $1 \le i, j \le N$ as in the proof of Lemma 3.8 or Lemma 3.9. Here δ_{ij} is Kronecker's delta. Let E' be the vector space spanned by e_i'. Since the support of all the e_i' are contained in the compact support of $\rho_{a,\epsilon}$, Assumption 3.5 implies that

$$\left| \mu_i \mu_j \delta_{ij} - \int_M (D_t e_i', D_t e_j') \right| < \epsilon_0.$$

for every t sufficiently closed to 0. Let E' be the vector space spanned by e_i'. Then if ϵ_0 is sufficiently small, each element s' of E' satisfies $||D_t s'||_M^2 \leq \lambda_1 ||s'||_M^2$. Corollary 3.15 implies $\dim E' \leq \dim \oplus_{\lambda<\lambda_1} E_\lambda(D_t)$. It is easy to check that the above inequality holds for each degree of $\mathbb{Z}/2$. □

Proof of Theorem 3.6. From Proposition 3.14 there is $0 < \lambda_1 < \lambda_0$ such that $E_{\lambda_1} = 0$. Then Corollary 3.18 and Proposition 3.19 imply

$$\dim \oplus_{\lambda\leq\lambda_1} E_\lambda(D_0) = \lim_{t\to 0} \dim \oplus_{\lambda\leq\lambda_1} E_\lambda(D_t)$$

and the equality holds for each degree, from which the claim follows. □

3.5. **Gluing formula.** In this subsection we generalize Theorem 3.6. We first need to generalize Assumption 3.5.

Let (M, W, σ, D, V) be as in Section 3.1. For $i = 1, 2, \ldots$ we further suppose that the data $(M_i, W_i, \sigma_i, D_i, V_i)$ satisfies the setting in Section 3.1.

Suppose M has a sequence of compact open subsets $\{K_i\}$ $(i = 1, 2, \ldots)$ satisfying $K_1 \cup V = M$, $K_i \subset int(K_{i+1})$ and $M = \cup_i K_i$. Suppose there exists an isometric open embedding $\iota_i : int(K_i) \to M_i$ and an isomorphism $\widetilde{\iota}_i : W|_{int(K_i)} \cong \iota_i^* W_i$ as $\mathbb{Z}/2$-graded Hermitian vector bundle over $int(K_i)$ for each i.

If s is a compactly supported section of W, then $\widetilde{\iota}_i s|_{int(K_i)}$ makes sense as a compactly supported section of W_i for large i with $supp\, s \subset int(K_i)$ by extending as 0 outside $\iota_i(supp\, s)$. We simply write $\widetilde{\iota}_i s$ for this section on W_i for large i.

If K is a compact subset of M and s_i is a section of W_i for large i with $K \subset int(K_i)$, then $\widetilde{\iota}_i^{-1}(s_i|_{\iota(K_i)})|_K$ is a section of $W|_K$. We simply write $\widetilde{\iota}_i^{-1} s_i|_K$ for this section.

Then $s \mapsto \{\widetilde{\iota}_i^{-1}(D_i \widetilde{\iota}_i s)|_K\}$ is a differential operator on K for large i. We write $\iota_i^* D_i$ for this operator on K.

Assumption 3.20.
(1) $\iota_i(M \smallsetminus V) = M_i \smallsetminus V_i$.
(2) The data $(M_i, W_i, \sigma_i, D_i, V_i)$ satisfy Assumption 3.1 with the same constant λ_0 for (M, W, σ, D, V).
(3) On each compact subset K of M the coefficients of the differential operator $\iota_i^* D_i$ is C^∞-convergent to those of D on K as $i \to \infty$.

We do not assume that the propagation speed of $\{D_i\}$ is uniformly bounded with respect to i.

Theorem 3.21. *Under Assumption 3.20,* ind D_i *is equal to* ind D *for large* i.

Proof. The most of the arguments in Sections 3.3 and 3.4 go through.

The statement and proof of Proposition 3.13 is straightforwardly generalized with replacement of D_{t_i} by D_i.

The statement and proof of Proposition 3.19 is also straightforwardly generalized.

To generalize Proposition 3.16 we need the construction in Remark 3.17. As for the statement we replace D_{t_i} with D_i, and let $E^{(i)}$ be a $\mathbb{Z}/2$ graded subspace of $L^2(M_i, W_i)$. As for the proof take $e_k^{(i')}$ with support contained in $int(K_i)$. Let $s_{i'}$ be an element of $E^{(i')}$ with $||s_{i'}||_{M_i} = 1$ which is L^2-orthogonal to all $\widetilde{\iota}_i e_k^{(i')}$ $(1 \leq k \leq N)$. Then the rest of the proof remain valid.

Then the argument of Section 3.4 can be straightforwardly generalized to show Theorem 3.21. □

When M has two (or more) connected components while each M_i' is connected, Theorem 3.21 is regarded as a gluing formula of index as explained below.

Proposition 3.22. *Let* (M, W, σ, D, V) *be a data satisfying Assumption 3.1. Suppose* M *is the disjoint union of* M' *and* M''*. Let* (W', σ', D', V') *and* $(W'', \sigma'', D'', V'')$ *be the restrictions of* (W, σ, D, V) *to* M' *and* M'' *respectively. Then we have*

$$\text{ind } D_i = \text{ind } D' + \text{ind } D'',$$

for sufficiently large i*.*

Proof. Since $\text{ind } D = \text{ind } D' + \text{ind } D''$, Theorem 3.21 implies the required gluing formula. □

The following vanishing lemma follows from the partial integration formula Lemma 3.8 and the second inequality in Assumption 3.1.

Lemma 3.23. *Let* (M, W, σ, D, V) *be a data satisfying Assumption 3.1. If* $M = V$*, then we have* $\ker D \cap L^2(W) = 0$.

Using the gluing formula Proposition 3.22 and the above Lemma 3.23, we have the following excision formula of index.

Proposition 3.24. *Let* (M, W, σ, D, V) *be a data satisfying Assumption 3.1. Suppose* M *is the disjoint union of* M' *and* M''*, and* M'' *is contained in* V*. Let* (W', σ', D', V') *be the restriction of* (W, σ, D, V) *to* M'*. Then we have*

$$\text{ind } D_i = \text{ind } D'$$

for sufficiently large i*.*

3.6. **Product formula.** Following Atiyah and Singer [1], we formulate a product formula for elliptic operators. Except that we need Lemma 3.8 for partial integration on complete Riemannian manifolds, the argument is exactly the same as in [1]. The main purpose of this subsection is to formulate Assumption 3.26 below, which is crucial for the case of complete manifolds. To apply the product formula it is necessary to check the assumption for specific operators, which is our another task and is carried out in Section 5.

For $k = 0, 1$ let M_k be a complete Riemannian manifold, W_k a $\mathbb{Z}/2$-graded Hermitian vector bundle over M_k, and $D_k : \Gamma(W_k) \to \Gamma(W_k)$ a degree-one formally self-adjoint order-one elliptic operator with principal symbol σ_k.

Let G be a compact Lie group and $P \to M_0$ a principal G-bundle. Suppose G acts on M_1 isometrically, W_1 is G-equivariant $\mathbb{Z}/2$-graded Hermitian vector bundle, and D_k is a G-invariant operator.

Then $M = P \times_G M_1$ is a fiber bundle over M_0 with fiber M_1. We write $\pi : M \to M_0$ for the projection map. Let $\widetilde{W_0}$ and $\widetilde{W_1}$ be the vector bundles over M defined by $\widetilde{W_0} = \pi^* W_0$ and $\widetilde{W_1} = P \times_G W_1$, and we put $W = \widetilde{W_0} \otimes \widetilde{W_1}$.

We would like to lift D_0 and D_1 as operators on W. The lift of D_1 is given straightforward: Defining the operator $\widetilde{D}_1$ on $\Gamma(\widetilde{W_0} \otimes \widetilde{W_1})$ by $\epsilon \otimes D_1$ on each fiber of $\pi : M \to M_0$, where $\epsilon : W_0 \to W_0$ is equal to $+id$ on the degree 0 part of W_0, and to $-\text{id}$ on the degree 1 part of W_1.

We next construct $\widetilde{D}_0 : \Gamma(W) \to \Gamma(W)$. Let $\{V_\alpha\}$ be an open covering of M_0 and $\{\rho_\alpha^2\}$ a partition of unity. Suppose we have local trivializations $P|_{V_\alpha} \cong V_\alpha \times G$ with transition functions $g_{\alpha\beta}$. Using the local trivialization on V_α we have the identifications

$\pi^{-1}(V_\alpha)\cong V_\alpha\times M_1$ and $W|_{\pi^{-1}(V_\alpha)}\cong W_0|_{V_\alpha}\times W_1$. Let $\widetilde{D}_{0,\alpha}$ be the operator on $W|_{\pi^{-1}(V_\alpha)}$ defined by D_0 using the product structure. We put $\widetilde{D}_0:=\sum_\alpha\rho_\alpha\widetilde{D}_{0,\alpha}\rho_\alpha$.

Lemma 3.25. $\widetilde{D}_0\widetilde{D}_1+\widetilde{D}_1\widetilde{D}_0=0$.

Proof. It follows from $\widetilde{D}_1\widetilde{D}_{0,\alpha}+\widetilde{D}_{0,\alpha}\widetilde{D}_1=0$ and $\widetilde{D}_1\rho_\alpha-\rho_\alpha\widetilde{D}_1=0$. ☐

Let R be a G-invariant $\mathbb{Z}/2$-graded finite dimensional subspace of $\Gamma(M_1,W_1)$, and $\widetilde{R}$ the fiber bundle $P\times_G R$ over M_0. Then we have an embedding

$$\Gamma(M_0,W_0\otimes\widetilde{R})\to\Gamma(M,\widetilde{W}) \tag{10}$$

which is preserved by the action of $\widetilde{D}_0$. Let $\widetilde{D}_R$ be the restriction of $\widetilde{D}_0$ on $\Gamma(M_0,W_0\otimes\widetilde{R})$. Then $\widetilde{D}_R$ is a differential operator on $W_0\otimes\widetilde{R}$ with principal symbol $\sigma_0\otimes\mathrm{id}_{\widetilde{R}}$.

Assumption 3.26.
1. D_0 has finite propagation speed, i.e., σ_0 is L^∞-bounded.
2. The data (M_1,W_1,D_1) satisfies Assumption 3.1.
3. $R=E_0(D_1)$.
4. The data $(M,W,\widetilde{D}_0+\widetilde{D}_1)$ satisfies Assumption 3.1.

We do not assume the second condition of Assumption 3.1 for the data (M_0,W_0,D_0).

Recall that $\widetilde{D}_R$ is given by $\widetilde{D}_0$ via the embedding (10). Since $\widetilde{D}_1=0$ on the image of the embedding, Assumption 3.26 implies that the data $(M_0,W_0\otimes\widetilde{R},\widetilde{D}_R$ satisfies Assumption 3.1 as well.

Theorem 3.27 (product formula)**.** $\mathrm{ind}(\widetilde{D}_0+\widetilde{D}_1)=\mathrm{ind}\,\widetilde{D}_R$

Proof. We show that the embedding (10) gives the isomorphism $E_0(\widetilde{D}_R)\cong E_0(\widetilde{D}_0+\widetilde{D}_1)$. If s is in the image of $E_0(D_R)$, then the construction of D_R implies that s is obviously in $E_0(\widetilde{D}_0+\widetilde{D}_1)$. From Lemma 3.8and Lemma 3.25 if s is an element of $E_0(\widetilde{D}_0+\widetilde{D}_1)$ we have

$$0=\int_M((\widetilde{D}_0+\widetilde{D}_1)^2s,s)=\int_M(\widetilde{D}_0^2s+\widetilde{D}_1^2s,s)=||\widetilde{D}_0s||^2+||\widetilde{D}_1s||^2,$$

i.e., $\widetilde{D}_0s=\widetilde{D}_1s=0$. In particular $\widetilde{D}_1s=0$ implies that s is in the image of the embedding (10). Moreover $\widetilde{D}_0s=0$ implies s is in the image of $E_0(D_R)$. ☐

4. Local index

In this section we first define a class of Riemannian manifolds, and consider compatible fibrations and compatible systems on them which satisfy some natural conditions. For such a class we will define the local index of a strongly acyclic compatible system in this section and prove the product formula in the next section.

Definition 4.1. Let M be a Riemannian manifold. If there exists an open subset V of M which satisfies the following properties, then we call M a *manifold with the generalized cylindrical end* V.

1. The complement $M\smallsetminus V$ is compact.
2. There exists a finite open covering $\{V_\beta\}_{\beta\in B}$ of V.
3. There exist families of Riemannian manifolds $\{N_\beta\}_{\beta\in B}$ and non-negative integers $\{m_\beta\}_{\beta\in B}$ such that each N_β is an open subset of a closed Riemannian manifold and each V_β is isometric to $N_\beta\times\mathbb{R}_{>0}^{m_\beta}$, where $\mathbb{R}_{>0}$ is the set of positive real numbers.

We identify V_β with $N_\beta \times \mathbb{R}^{m_\beta}_{>0}$ and write $V = \cup_\beta N_\beta \times \mathbb{R}^{m_\beta}_{>0}$.

Let M be a manifold with the generalized cylindrical end $V = \cup_{\beta\in B} N_\beta \times \mathbb{R}^{m_\beta}_{>0}$. Let $\{\pi_\alpha, V_\alpha\}_{\alpha\in A}$ be a compatible fibration on V. Hereafter we take and fix a relatively compact open neighborhood V_∞ of $M \smallsetminus V$ and put $\widetilde{A} := A \cup \{\infty\}$. Let $\{\rho^2_\alpha\}_{\alpha\in\widetilde{A}}$ be an admissible partition of unity of $M = \cup_{\alpha\in\widetilde{A}} V_\alpha$. Here V_∞ has the fiber bundle structure whose fiber consists of a single point. Let $\{W, D_\alpha\}_{\alpha\in A}$ be a compatible system on V. In this section we consider the above data with the following *translationally invariance* conditions on the end.

Condition (TI)

1. There is a bijective map between A and B.
2. Each V_α is isometric to $N_\alpha \times \mathbb{R}^{m_\alpha}_{>0}$ under the identification in (1).
3. Each N_α has a fiber bundle structure $N_\alpha \to U_{N_\alpha}$ such that the fiber bundle structure $V_\alpha \to U_\alpha$ can be identified with the induced fiber bundle structure $N_\alpha \times \mathbb{R}^{m_\alpha}_{>0} \to U_{N_\alpha} \times \mathbb{R}^{m_\alpha}_{>0}$.
4. For each $\alpha \in A$, the $\mathbb{Z}/2$-graded Clifford module bundle $W|_{V_\alpha}$ and the operator D_α are invariant under the translation action of the $\mathbb{R}^{m_\alpha}_{>0}$-direction.
5. For each $\alpha \in A$, there exist an open subset N'_α of N_α and a positive real number ϵ_α such that $N'_\alpha \Subset N_\alpha$ and $M = V_\infty \cup (\cup_{\alpha\in A} N'_\alpha \times \mathbb{R}^{m_\alpha}_{>\epsilon_\alpha})$, where $\mathbb{R}_{>\epsilon_\alpha}$ is the set of real numbers greater than ϵ_α.
6. For each $\alpha \in A$ there exists a compactly supported function on N'_α such that $\rho_\alpha|_{N'_\alpha \times \mathbb{R}^{m_\alpha}_{>\epsilon_\alpha}}$ is the pull back of it via the natural projection $N'_\alpha \times \mathbb{R}^{m_\alpha}_{>\epsilon_\alpha} \to N'_\alpha$.

A typical example of a data with Condition (TI) is a data on a manifold with cylindrical end $V = N \times \mathbb{R}_+$ which is invariant under the translation on the $\mathbb{R}_+$-direction. We can define a product of two data with Condition (TI) in an appropriate and obvious way. The product also satisfies the same condition, which is crucial in the proof of the product formula. See Section 5 for details.

We first show the following technical lemma for the above data which is used to show Theorem 4.7.

Lemma 4.2. *Suppose that A is non-empty and V is connected. There exists an admissible partition of unity $\{\chi^2_\alpha\}_{\alpha\in\widetilde{A}}$ which satisfies the condition (5),(6) in Condition (TI) and* $\mathrm{supp}(\chi_\alpha) \Subset \mathrm{supp}(\rho_\alpha)$ *for each $\alpha \in A$.*

Proof. Since A is a finite set it is enough to show that when we fix $\alpha \in A$, there exists an admissible partition of unity $\{\chi^2_{\alpha'}\}_{\alpha'\in\widetilde{A}}$ of $M = \cup_{\alpha'\in\widetilde{A}} V_{\alpha'}$ which satisfies the condition (5) and (6) in Condition (TI), $\mathrm{supp}(\chi_\alpha) \Subset \mathrm{supp}(\rho_\alpha)$ and $\mathrm{supp}(\chi_{\alpha'}) = \mathrm{supp}(\rho_{\alpha'})$ for all $\alpha' \in \widetilde{A} \smallsetminus \{\alpha\}$.

We first take and fix a non-decreasing function $\varphi : \mathbb{R}_{\geq 0} \to \mathbb{R}$ such that

$$\varphi(r) = \begin{cases} 0 & (0 \leq r \leq 1/2) \\ r & (1 \leq r), \end{cases}$$

and $\mathrm{supp}(\varphi) = [1/2, \infty)$. We define $\hat{\rho}_\alpha : M \to \mathbb{R}$ by the composition $\hat{\rho}_\alpha := \varphi \circ \rho_\alpha$. Then, by definition, we have $\mathrm{supp}(\hat{\rho}_\alpha) \Subset \mathrm{supp}(\rho_\alpha)$ and

$$\hat{\rho}_\alpha(x) + \sum_{\alpha'\neq\alpha} \rho_{\alpha'}(x) > 0$$

for all $x \in M$. By normalizing the family of functions $\{\hat{\rho}_\alpha\} \cup \{\rho_{\alpha'}\}_{\alpha' \neq \alpha}$ we obtain the required admissible partition of unity $\{\chi^2_{\alpha'}\}_{\alpha' \in \widetilde{A}}$. ☐

4.1. **Vanishing theorem.** Let M be a Riemannian manifold and V an open subset of M. Let $\{\pi_\alpha : V_\alpha \to U_\alpha\}_{\alpha \in A}$ be a compatible fibration and $\{W, D_\alpha\}_{\alpha \in A}$ a compatible system on V. Let $\{\rho^2_\alpha\}_{\alpha \in \widetilde{A}}$ be an admissible partition of unity. In this subsection we assume the following three conditions.

- M has a generalized cylindrical end V'.
- $\{\pi_\alpha : V_\alpha \to U_\alpha, W, D_\alpha, \rho_\alpha\}$ satisfies Condition (TI).
- $\{W, D_\alpha\}_{\alpha \in A}$ is strongly acyclic.

For each $\alpha \in A$ we put $D'_\alpha := \rho_\alpha D_\alpha \rho_\alpha$. Take any Dirac-type operator D on $\Gamma(W)$ which is translationally invariant on the end and a positive real number t. We define the operator acting on $\Gamma(W)$ by $D_t := D + t\sum_{\alpha \in A} D'_\alpha$.

Remark 4.3. Note that D_α and ρ_β commute each other because ρ_β is a pull-back of a function on U_β and D_α contains only the derivatives along fibers.

Proposition 4.4. *D_t is elliptic for all $t \geq 0$.*

Proof. With the identification of tangent bundles and cotangent bundles via the Riemannian metric, the principal symbol σ_t of D_t is written as

$$\sigma_t = c \circ (\mathrm{id}_{TM} + t\sum_\alpha \rho_\alpha p_\alpha \rho_\alpha),$$

where $p_\alpha \colon TV_\alpha \to T[\pi_\alpha]$ is the orthogonal projection. We put

$$S := \mathrm{id}_{TM} + t\sum_\alpha \rho_\alpha p_\alpha \rho_\alpha.$$

S is a symmetric operator. In order to prove the proposition it is sufficient to show that for each $x \in M$ $S_x \colon T_xM \to T_xM$ is a positive operator. By definition, for $u \in T_xM$ we have

$$(S_x u, u) = |u|^2 + t\sum_\alpha |\rho_\alpha(x) p_\alpha(u)|^2 \geq 0,$$

where we think $p_\alpha(u) = 0$ if $x \notin V_\alpha$. Moreover, $(S_x u, u) = 0$ if and only if $u = 0$. This implies S_x is positive. ☐

We show the following fundamental lemma in our argument.

Lemma 4.5. *For each α the anti-commutator $DD'_\alpha + D'_\alpha D$ is a differential operator along fibers of π_α of order at most 2.*

Proof. Recall that, for each α the principal symbol of horizontal direction D^{H_α} of D with respect to π_α anti-commutes not only with the symbol of D'_α, but also with the whole operator D'_α. The statement follows from this property. It is straightforward to check it using local description. Instead of giving the detail of the local calculation, however, we here give an alternative formal explanation for the above lemma. For $b \in U_\alpha$ let $\mathcal{W}_b$ be the sections of the restriction of W on the fiber $\pi_\alpha^{-1}(b)$. Then $\mathcal{W} = \coprod \mathcal{W}_b$ is formally an infinite dimensional vector bundle over U_α. We can regard D'_α as an endmorphism on $\mathcal{W}$. Then D'_α is a order-zero differential operator on $\mathcal{W}$ whose principal symbol is equal to D'_α itself. Then, as a differential operator on $\mathcal{W}$, the anti-commutator $D'_\alpha D^{H_\alpha} + D^{H_\alpha} D'_\alpha$ is an (at most) order-one operator whose principal symbol is given by the anti-commutator between the Clifford action by TU_α and D'_α. This principal symbol vanishes, which implies that the anti-commutator is order-zero as a differential operator on $\mathcal{W}$, i.e., it does not contain derivatives of U_α-direction. ☐

For an operator appearing in the above lemma we have the following a priori estimate.

Lemma 4.6. *For each fiber F of π_α and arbitrary differential operator Q of order at most 2 along F, there exists a constant C_Q such that the inequality*

$$\left|\int_F (s_F, Qs_F)\right| \leq C_Q \int_F |D_\alpha s_F|^2$$

holds for all sections s_F.

The following vanishing theorem is a main theorem in this subsection.

Theorem 4.7. *Let M be a Riemannian manifold with the generalized cylindrical end V. Suppose that M is equipped with the data $\{\pi_\alpha : V_\alpha \to U_\alpha, W, D_\alpha, \rho_\alpha\}$ which satisfies Condition (TI). Let D be a Dirac-type operator acting on $\Gamma(W)$ which is translationally invariant on the end. Put $D'_\alpha := \rho_\alpha D_\alpha \rho_\alpha$ and $D_t := D + t\sum_\alpha D'_\alpha$ for a positive number $t > 0$. Then the space of L^2-solutions of the equation $D_t s = 0$ is trivial for all $t \gg 1$.*

To show this theorem we make several preparations. Let $\{\chi_\alpha^2\}_{\alpha\in\widetilde{A}}$ be an admissible partition of unity constructed in Lemma 4.2 and put $K_\alpha :=$ *supp* χ_α.

Lemma 4.8.
1. *D'_α is an elliptic operator on each fiber of π_α which is contained in K_α.*
2. *If Q is a differential operator along fibers of order at most 2, then there exists a constant C_Q such that for each section s_α satisfying supp $s_\alpha \subset K_\alpha$ and $D'_\alpha s_\alpha \in L^2(W)$ we have an estimate*

$$\left|\int_M (s_\alpha, Qs_\alpha)\right| \leq C_Q \int_M |D'_\alpha s_\alpha|^2.$$

Proof. The first statement follows from the fact that ρ_α takes positive values on K_α. Note that since the data satisfies Condition (TI) we can choose the constants C_Q in Lemma 4.6 uniformly for all fibers of π_α contained in K_α. The second statement follows from this fact. ☐

Lemma 4.9. *There exists an operator Z which does not contain any differential terms and satisfying*

$$D_t^2 = \sum_\alpha \chi_\alpha D_t^2 \chi_\alpha + Z.$$

Moreover Z does not depend on t.

Proof. If χ is an admissible function then it commutes with D_α and hence we have $[D_t, \chi] = [D, \chi]$. Using this equality and the fact $[D, \chi]$ does not contain any differential operators we have

$$\begin{aligned}
[[D_t^2, \chi], \chi] &= [(D_t[D_t, \chi] + [D_t, \chi]D_t), \chi] \\
&= [(D_t[D, \chi] + [D, \chi]D_t), \chi] \\
&= [D_t, \chi][D, \chi] + [D, \chi][D_t, \chi] \\
&= 2[D, \chi]^2.
\end{aligned}$$

Put $\chi := \chi_\alpha$ and take summation for all α we have

$$2D_t^2 - 2\sum_\alpha \chi_\alpha D_t^2 \chi_\alpha = \sum_\alpha [[D_t^2, \chi], \chi] = 2\sum_\alpha [D, \chi_\alpha]^2.$$

Then $Z := \sum_\alpha [D, \chi_\alpha]^2$ is the required operator of order 0. □

Proposition 4.10. *If s_α is an L^2-bounded section of W such that Ds_α and $D'_\alpha s_\alpha$ are also L^2-bounded section for all $\alpha \in A$ and supp $s_\alpha \subset K_\alpha$, then we have the inequality*

$$\int_M |D_t s_\alpha|^2 \geq \left| \int_M (Zs_\alpha, s_\alpha) \right| + \int_M |s_\alpha|^2$$

for all $t \gg 1$.

We use the above proposition and lemmas to show Theorem 4.7 as follows.

Proof of Theorem 4.7 assuming Proposition 4.10. We take $t \gg 1$ so that the inequality in Proposition 4.10 holds. We first note that since M has a generalized cylindrical end and the data satisfies Condition (TI), an L^2-bounded section s which satisfies $D_t s = 0$ is an element in the Sobolev space $L^2_k(W)$ for arbitrary $k \in \mathbb{N}$ by the elliptic estimate, and hence $s_\alpha = \chi_\alpha s$ is. Moreover since D_t is translationally invariant it has the bounded extension $D_t : L^2_1(W) \to L^2(W)$. Then we can use the partial integration formula in Lemma 3.8 for $D_t s_\alpha$, and we have $s = 0$ as in the following;

$$\begin{aligned}
0 &= \int_M (D_t^2 s, s) \\
&= \sum_\alpha \int_M (\chi_\alpha D_t^2 \chi_\alpha s, s) + \int_M (Zs, s) \qquad \text{(Lemma 4.9.1)} \\
&= \sum_\alpha \left(\int_M |D_t s_\alpha|^2 + \int_M (Zs_\alpha, s_\alpha) \right) \qquad (s_\alpha := \chi_\alpha s) \\
&\geq \sum_\alpha \left(\int_M |(Zs_\alpha, s_\alpha)| + \int_M |s_\alpha^2| + \int_M (Zs_\alpha, s_\alpha) \right) \qquad \text{(Proposition 4.10)} \\
&\geq \sum_\alpha \int_M |s_\alpha^2| = \int_M |s|^2.
\end{aligned}$$

□

4.1.1. *Proof of Proposition 4.10.* For each fixed $\alpha \in A$, we can write D_t as

$$D_t = D_{\neq\alpha} + tD'_\alpha$$

on V_α, where we put

$$D_{\neq\alpha} := D + t\sum_{\beta\neq\alpha} D'_\beta.$$

Using these notations together with $Q_\alpha := DD'_\alpha + D'_\alpha D$ and $Q_{\beta\alpha} := D'_\beta D'_\alpha + D'_\alpha D'_\beta$, we have

$$D_t^2 = D_{\neq\alpha}^2 + tQ_\alpha + t^2 \sum_{\beta\neq\alpha} Q_{\beta\alpha} + t^2 D'^2_\alpha,$$

and since $\int_M (D^2_{\neq\alpha} s_\alpha, s_\alpha)$ and $\int_M (Q_{\beta\alpha} s_\alpha, s_\alpha)$ are non-negative for an L^2-section s_α satisfying the assumptions we also have

$$\int_M |D_t s_\alpha|^2 \geq t^2 \int_M |D'_\alpha s_\alpha|^2 - t \left| \int_M (Q_\alpha s_\alpha, s_\alpha) \right|.$$

From Lemma 4.5 Q_α is a differential operator along fibers of π_α of order at most 2. Then from Lemma 4.8.2 there exist a constant C' such that

$$\left| \int_M (Q_\alpha s_\alpha, s_\alpha) \right| \leq C' \int_M |D'_\alpha s_\alpha|^2.$$

Combining these inequalities we have

$$\int_M |D_t s_\alpha|^2 \geq (t^2 - C't) \int_M |D'_\alpha s_\alpha|^2.$$

On the other hand using Lemma 4.8.2 again there exists a constant C'' such that

$$\left| \int_M (Z s_\alpha, s_\alpha) \right| + \int_M |s_\alpha|^2 \leq C'' \int_M |D'_\alpha s_\alpha|^2,$$

and hence if we take $t \gg 1$ so that $t^2 - C't \geq C''$, then we have

$$\int_M |D_t s_\alpha|^2 \geq \left| \int_M (Z s_\alpha, s_\alpha) \right| + \int_M |s_\alpha|^2.$$

Note that since A is a finite set we may assume that C' and C'' do not depend on α, and we complete the proof.

4.2. **Definition of the local index - Generalized cylindrical end case.** In this subsection we give the definition of the local index of a strongly acyclic compatible system on a manifold M with the generalized cylindrical end V. Consider the data on V which satisfies the Condition (TI) and take any Dirac-type operator D acting on $\Gamma(W)$ which is translationally invariant on the end. We assume that the compatible system on V is strongly acyclic. As in the previous Subsection we put $D_t := D + t\sum_\alpha \rho_\alpha D_\alpha \rho_\alpha$.

Lemma 4.11. *If t is large enough so that the inequality in Proposition 4.10 holds, then D_t satisfies the Assumption 3.1.*

Proof. Since the principal symbol of D_t is given by a linear combination of the Clifford multiplication of TM and that of fiber directions of $\{\pi_\alpha\}_{\alpha\in A}$, (i) of Assumption 3.1 is satisfied. We show the condition (ii) of Assumption 3.1. Let s be a smooth compactly-supported section of W with *supp* $s \subset V$. Let $\{\chi^2_\alpha\}_{\alpha\in\widetilde{A}}$ be the admissible partition of unity constructed in Lemma 4.2. For each $s_\alpha := \chi_\alpha s$ we can apply Proposition 4.10, and hence, we have $||D_t s||^2_V \geq ||s||^2_V$ as in the same way in the proof of Theorem 4.7. □

Results in Section. 3 imply the following.

Proposition 4.12. *If t is large enough so that the inequality in Proposition 4.10 holds, then the space of L^2-solutions of $D_t s = 0$ is finite dimensional and its super-dimension is independent for $t \gg 1$ and any other continuous deformations of data.*

Definition 4.13. We define the local index $\mathrm{ind}(M, V, W)$ as the index of D_t in the sense of Section 3.

In the case of cylindrical end we have the following sum formula of local indices.

Lemma 4.14. *For $i=1,2$ let M_i be manifolds with cylindrical ends $V_i = N_i \times \mathbb{R}_{>0}$ and N_i^0 be connected components of N_i^0. Suppose that there is an isometry $\phi : N_1^0 \to N_2^0$, and for some $R>0$ the map $\phi : N_1^0 \times (0,R) \to N_2^0 \times (0,R)$ given by $(x,r) \mapsto (\phi(x), R-r)$ induces the isomorphism between the strongly acyclic compatible systems on them. Then we can glue $M_1 \smallsetminus (N_1^0 \times [R,\infty))$ and $M_2 \smallsetminus (N_2^0 \times [R,\infty))$ to obtain a new manifold $\hat{M}$ with cylindrical end $\hat{V} = \hat{N} \times (0,\infty)$ for $\hat{N} = (N_1 \smallsetminus N_1^0) \cup (N_2 \smallsetminus N_2^0)$, and we also have a Clifford module bundle $\hat{W}$ obtained by gluing W and W' on $N_1^0 \times (0,R) \cong N_2^0 \times (0,R)$. Then we have*

$$\mathrm{ind}(\hat{M}, \hat{V}, \hat{W}) = \mathrm{ind}(M_1, V_1, W_1) + \mathrm{ind}(M_2, V_2, W_2).$$

4.3. **Definition of the local index - general case.** Let V be an open subset of M such that $M \smallsetminus V$ is compact. Assume that V has a strongly acyclic compatible system. We would like to define the local index for such a general case. The way to define it is almost same in [4]. To verify the construction we have to check the following.

Proposition 4.15. *For given (M,V,W) and the strongly acyclic compatible system $(V_\alpha \overset{\pi_\alpha}{\to} U_\alpha, D_\alpha)$ on V we can deform them to (M', V', W') so that it has a cylindrical end with a translationally invariant strongly acyclic compatible system.*

To prove the proposition, it is enough to show the following lemma.

Lemma 4.16. *There exists a smooth admissible function $f : M \to \mathbb{R}$ and a regular value c of f such that $f^{-1}(-\infty, c]$ is a compact subset containing $M \smallsetminus V$.*

To show this lemma, we first note that the following.

Lemma 4.17. *Let $\{V''_\alpha\}_{\alpha \in A}$ be the open covering of M which appears in the 4th property in Definition 2.11. For any subset D of M, let $K(D)$ be*

$$K(D) = \bigcup_\alpha \pi_\alpha^{-1}\pi_\alpha(D \cap \overline{V''_\alpha}).$$

Let $h : M \to \mathbb{R}$ be a smooth function. Let J be a connected interval. If $K(D) \subset h^{-1}(J)$, then we have $D \subset I(h)^{-1}(J)$, where $I : C^\infty(M) \to C^\infty(M)$ is the averaging operation.

Proof. Use the fact $\pi_\alpha^{-1}\pi_\alpha(V''_\alpha) = V''_\alpha$ and the inequality in Definition 2.11. □

Proof of Lemma 4.16. Since each π_α is a proper map, the subset $K(M \smallsetminus V)$ defined in Lemma 4.17 is compact. Let $f_0 : M \to \mathbb{R}$ be the distance function from the compact subset $K(M \smallsetminus V)$. Take a real number $r > 0$ so that $f_0^{-1}[0,r]$ is a compact neighborhood of $K(M \smallsetminus V)$. Let $\epsilon > 0$ be a positive real number satisfying $2\epsilon < r$. Let $h : M \to \mathbb{R}$ be a smooth function such that $|f_0(x) - h(x)| < \epsilon$ for all $x \in M$, and we put $f := I(h)$. Let c be a regular value of f satisfying $\epsilon < c < r - \epsilon$. Then we have

$$K(M \smallsetminus V) = f_0^{-1}(0) \subset h^{-1}(-\epsilon, \epsilon).$$

By Lemma 4.17 we have

$$M \smallsetminus V \subset f^{-1}(-\epsilon, \epsilon) \subset f^{-1}(-\infty, c].$$

Suppose that $x \notin K(h^{-1}(-\infty, c])$. Then one has $K(\{x\}) \subset h^{-1}(c, \infty)$, and it implies $x \notin f^{-1}(-\infty, c]$ by Lemma 4.17. Then we have

$$f^{-1}(-\infty, c] \subset K(h^{-1}(-\infty, c]) \subset K(f_0^{-1}(-\infty, c+\epsilon]) \subset K(f_0^{-1}[0, r]).$$

In particular $f^{-1}(-\infty, c]$ is compact. □

Definition 4.18. We define the local index $\operatorname{ind}(M, V, W)$ to be the local index for the deformed data (M', V', W').

Note that the local index $\operatorname{ind}(M, V, W)$ is well-defined, i.e, it does not depend on various choice of the construction. It follows from the sum formula (Lemma 4.14) and Theorem 4.7 as in the same way in [4]. The well-definedness means the excision property of local index.

Theorem 4.19. *Let M be a Riemannian manifold and V an open subset such that $M \smallsetminus V$ is compact. Let W be a $\mathbb{Z}_2$-graded $Cl(TM)$-module bundle on M and suppose that the metric on V is a compatible metric and V has a strongly acyclic compatible system. Let M' be an open neighborhood of $M \smallsetminus V$ with $V \cap M'$ admissible. Then we have*

$$\operatorname{ind}(M, V, W) = \operatorname{ind}(M', V \cap M', W|_{M'}).$$

The local index $\operatorname{ind}(M, V, W)$ also has the following property.

Theorem 4.20. *Let M, V, and W be the data as in Theorem 4.19. Suppose that V' is an admissible open subset of V with $M \smallsetminus V'$ compact. Then we have*

$$\operatorname{ind}(M, V, W) = \operatorname{ind}(M, V', W).$$

Note that if M is closed, then $\operatorname{ind}(M, V, W)$ is equal to the index of the Dirac-type operator D because of the homotopy invariance of indices. Using the excision property, additivity for disjoint unions and vanishing theorems we have the localization theorem.

Theorem 4.21. *Let M be a closed Riemannian manifold and V an open subset. Let W be a $Cl(TM)$-module bundle on M and D a Dirac-type operator acting on $\Gamma(W)$. Suppose that V has a strongly acyclic compatible system. Let $\cup_{i=1}^{N} V_i$ be an open neighborhood of $M \smallsetminus V$ such that $V_i \cap V_j = \emptyset$ if $i \neq j$. Then we have the following equality.*

$$\operatorname{ind} D = \sum_{i=1}^{N} \operatorname{ind}(V_i, V_i \cap V, W|_{V_i \cap V}).$$

Remark 4.22. The arguments in this section are valid in orbifold category.

5. Product formula of local indices

In this section we formulate the product of acyclic compatible systems. Once we have an appropriate formulation of the product, then we obtain the product formula of local indices of the strongly acyclic compatible systems by results in Section 3.

5.1. **Product of compatible fibrations.**

In this subsection we formulate a product of compatible fibrations. The product is defined for the following collection of data for $i = 0, 1$ which satisfy the Assumption 5.1.

1. M_i : a manifold.
2. V_i : an open set of M_i.
3. $\{\pi_{i,\alpha} : V_{i,\alpha} \to U_{i,\alpha},\ U_{i,\alpha\beta} \mid \alpha, \beta \in A_i\}$: a compatible fibration on V_i.
4. G : a compact Lie group which acts smoothly on M_1.
5. $\pi_P : P \to M_0$: a principal G-bundle over M_0.

Assumption 5.1.

1. V_1 is G-invariant and the fibrations $V_{1,\alpha} \to U_{1,\alpha}$, $V_{1,\alpha} \cap V_{1,\beta} \to U_{1,\alpha\beta}$ and $U_{1,\alpha\beta} \to \pi_{1,\alpha}(V_{1,\alpha} \cap V_{1,\beta})$ are G-equivariant fiber bundles for all $\alpha, \beta \in A_1$.

(2) there exist principal G-bundles $P_\alpha \to U_{0,\alpha}$, $P_{\alpha\beta} \to U_{0,\alpha\beta}$ and bundle maps $P|_{V_{0,\alpha}\cap V_{0,\beta}} \to P_\alpha|_{\pi_{0,\alpha}(V_{0,\alpha}\cap V_{0,\beta})}$, $P|_{V_{0,\alpha}\cap V_{0,\beta}} \to P_{\alpha\beta}$, $P_{\alpha\beta} \to P_\alpha|_{\pi_{0,\alpha}(V_{0,\alpha}\cap V_{0,\beta})}$ for all $\alpha, \beta \in A_0$ such that the following diagrams commute;

$$\begin{tikzcd}
 & P|_{V_{0,\alpha}\cap V_{0,\beta}} \arrow[dl] \arrow[d] \arrow[dr] & \\
P_\alpha|_{\pi_{0,\alpha}(V_{0,\alpha}\cap V_{0,\beta})} \arrow[d] & V_{0,\alpha}\cap V_{0,\beta} \arrow[dl, "\pi_{0,\alpha}"'] \arrow[dr, "\pi_{\alpha\beta}"] & P_{\alpha\beta} \arrow[d] \\
\pi_{0,\alpha}(V_{0,\alpha}\cap V_{0,\beta}) & & U_{0,\alpha\beta} \arrow[ll, "p^{\alpha}_{\alpha\beta}"']
\end{tikzcd}$$

$$\begin{tikzcd}
 & P|_{V_{0,\alpha}\cap V_{0,\beta}} \arrow[ddl] \arrow[d] \arrow[ddr] & \\
 & P_{\alpha\beta} \arrow[dl] \arrow[dr] & \\
P_\alpha|_{\pi_{0,\alpha}(V_{0,\alpha}\cap V_{0,\beta})} & & P_\beta|_{\pi_{0,\beta}(V_{0,\alpha}\cap V_{0,\beta})}
\end{tikzcd}$$

For later convenience we take an open neighborhood $V_{i,\infty}$ of $M_i \smallsetminus V_i$ and consider the trivial fiber bundle structure $\pi_{i,\infty} : V_{i,\infty} \to V_{i,\infty}$. In other words we consider a compatible fibration $\{\pi_{i,\alpha} : V_{i,\alpha} \to U_{i,\alpha} \mid \alpha \in A_i \cup \{\infty\}\}$ on $M_i = V_{i,\infty} \cup (\cup_\alpha V_{i,\alpha})$. Let M be the quotient manifold by the diagonal action of G on $P \times M_1$. Then the natural map $\pi : M \to M_0$ is a fiber bundle whose fiber is equal to M_1. To define a structure of compatible fibration on M we first prepare several notations for $i = 0, 1$.

- $\widetilde{A}_i := A_i \cup \{\infty\}$.
- $\widetilde{A} := \widetilde{A}_0 \times \widetilde{A}_1$.
- $A := \widetilde{A} \smallsetminus (\infty, \infty)$.
- $V_{\alpha_0,\alpha_1} := P|_{V_{0,\alpha_0}} \times_G V_{1,\alpha_1}$ for $\alpha \in \widetilde{A}_i$.
- $U_{\alpha_0,\alpha_1} := P_{\alpha_0} \times_G U_{1,\alpha_1}$ for $\alpha \in \widetilde{A}_i$.
- $U_{(\alpha_0,\alpha_1)(\beta_0,\beta_1)} := P_{\alpha_0\beta_0} \times_G U_{1,\alpha_1\beta_1}$.
- $V := \bigcup_{(\alpha_0,\alpha_1)\in A} V_{\alpha_0,\alpha_1}$.

Then we have the following.

Proposition 5.2. *A collection of data*

$$\{\pi_{\alpha_0,\alpha_1} : V_{\alpha_0,\alpha_1} \to U_{\alpha_0,\alpha_1},\ U_{(\alpha_0,\alpha_1)(\beta_0,\beta_1)} \mid (\alpha_0,\alpha_1), (\beta_0,\beta_1) \in A\}$$

is a compatible fibration on $V = \bigcup_{(\alpha_0,\alpha_1)\in A} V_{\alpha_0,\alpha_1}$.

5.2. **Product of acyclic compatible systems.** In this subsection we define a product of acyclic compatible systems. To define the product we consider the following data together with the data 1, 2, 3, 4 and 5 in Subsection 5.1.

(6) a compatible Riemannian metric on M_i.

(7) W_i : a compatible $Cl(TM_i)$-module bundle over M_i.

(8) $\{D_{i,\alpha} \mid \alpha \in A_i\}$: strongly acyclic compatible system over $V_i = \cup_{\alpha\in A_i} V_{i,\alpha}$.

Together with Assumption 5.1 we assume the following.

Assumption 5.3. The metric on M_1 is G-invariant, and $W_1 \to M_1$ and $\{D_{1,\alpha}\}$ are G-equivariant.

From the Assumption 5.1 (2). the restrictions of P at each fibers of $\pi_{0,\alpha}$ are trivial. Moreover we have the following.

Lemma 5.4. *There exists a connection on P which is trivial flat over each fibers of $\pi_{0,\alpha}$ for all $\alpha \in A_0$.*

Proof. Take connections $\bar{\nabla}_\alpha$ for each $P_\alpha \to U_{0,\alpha}$. Let ∇_α be the pull-back connections of them to $P|_{V_{0,\alpha}} \to V_{0,\alpha}$ by $\pi_{0,\alpha}$. Define a connection ∇ on P by patching $\{\nabla_\alpha\}_\alpha$ by an admissible partition of unity $\{\rho_\alpha^2\}_\alpha$, which satisfies the required property. ☐

Using this connection on P and the compatible metric on M_0 we have the metric on P, and hence the metric on M. Moreover since the connection is trivial along fibers of $\{\pi_{0,\alpha}\}$ it induces a family of connections of $\{P_\alpha\}$ and $\{P_{\alpha\beta}\}$, and hence a family of metrics on them. Combining them with the G-invariant compatible metric on M_1 we have a family of metrics of $\{T[\pi_{\alpha_0\alpha_1}]\}$, $\{TU_{\alpha_0\alpha_1}\}$ and so on. It defines a compatible metric of a compatible fibration $M = \cup_{(\alpha_0,\alpha_1)} V_{\alpha_0,\alpha_1}$. We put $\widetilde{W_0} := \pi^* W_0 = \pi_P^* W_0 \times_G M_1$, $\widetilde{W_1} := P \times_G W_1$ and $W := \widetilde{W_0} \otimes \widetilde{W_1}$. Then $W \to M$ is a compatible Clifford module bundle of $M = \cup_{(\alpha_0,\alpha_1)} V_{\alpha_0,\alpha_1}$ with respect to the above induced compatible metric. Now we define differential operators $\widetilde{D}_{0,\alpha_0}$ and $\widetilde{D}_{1,\alpha_1}$ for each $\alpha_0 \in A_0$ and $\alpha_1 \in A_1$ which act on $\Gamma(\widetilde{W_0}|_{V_{\alpha_0,\alpha_1}})$ and $\Gamma(\widetilde{W_1}|_{V_{\alpha_0,\alpha_1}})$ respectively. The operator $\widetilde{D}_{1,\alpha_1}$ is the one induced from the G-equivariant operator D_{1,α_1} on $\Gamma(W_1|_{V_{1,\alpha_1}})$. On the other hand $\widetilde{D}_{0,\alpha_0}$ is the operator defined as follows: Since D_{0,α_0} is a differential operator along fibers of π_{0,α_0} and P is trivial at each fiber of π_{0,α_0}, we can define the operator acting on the restriction $\Gamma(\pi_P^* W_0|_{\mathrm{fiber}} \times_G M_1)$ using $D_{0,\alpha_0}|_{\mathrm{fiber}}$ and a trivialization of $P|_{\mathrm{fiber}}$. Since such operators along fibers do not depend on trivialization we have a differential operator $\widetilde{D}_{0,\alpha_0}$ acting on $\Gamma(\widetilde{W_0}|_{V_{\alpha_0,\alpha_1}})$. Using these operators we define an operator acting on $\Gamma(W|_V)$ by $D_{\alpha_0,\alpha_1} := \widetilde{D}_{0,\alpha_0} \otimes \mathrm{id}_{\widetilde{W_1}} + \epsilon_{\widetilde{W_0}} \otimes \widetilde{D}_{1,\alpha_1}$, where $\epsilon_{\widetilde{W_0}}$ is a map on $\widetilde{W_0}$ defined by $\epsilon_{\widetilde{W_0}}(v) := (-1)^{\deg v} v$. For later convenience we put $\widetilde{D}_{\alpha_0,\infty} = \widetilde{D}_{\infty,\alpha_1} = 0$.

Proposition 5.5. *A collection of differential operators $\{D_{\alpha_0,\alpha_1} \mid (\alpha_0,\alpha_1) \in A\}$ is a strongly acyclic compatible system on $(V, W|_V)$.*

Proof. Since $\widetilde{D}_{0,\alpha_0} \otimes \mathrm{id}_{\widetilde{W_1}}$ and $\epsilon_{\widetilde{W_0}} \otimes \widetilde{D}_{1,\alpha_1}$ anti-commute each other we have

$$\left(\sum t_{\alpha_0,\alpha_1} D_{\alpha_0,\alpha_1}\right)^2 = \left(\sum_{\alpha_0}\left(\sum_{\alpha_1} t_{\alpha_0,\alpha_1}\right)\widetilde{D}_{0,\alpha_0}\right)^2 \otimes \mathrm{id}_{\widetilde{W_1}} + \mathrm{id}_{\widetilde{W_0}} \otimes \left(\sum_{\alpha_1}\left(\sum_{\alpha_0} t_{\alpha_0,\alpha_1}\right)\widetilde{D}_{1,\alpha_1}\right)^2$$

for any family of non-negative numbers (t_{α_0,α_1}), and the equality among anti-commutators

$$\{D_{\alpha_0,\alpha_1}, D_{\alpha_0',\alpha_1'}\} = \{\widetilde{D}_{0,\alpha_0}, \widetilde{D}_{0,\alpha_0'}\} \otimes \mathrm{id}_{\widetilde{W_1}} + \mathrm{id}_{\widetilde{W_0}} \otimes \{\widetilde{D}_{1,\alpha_1}, \widetilde{D}_{1,\alpha_1'}\}.$$

These equalities imply that if $\{D_{i,\alpha_i}\}_{\alpha_i}$ are strongly acyclic compatible systems, then $\{D_{\alpha_0,\alpha_1}\}_{(\alpha_0,\alpha_1)}$ is so. ☐

5.3. **Product formula.** To apply results in Subsection 4.1 and 4.2, we have to deform the end of M_i and M as in the following way: we can deform M_i into $\hat{M}_i$ so that $\hat{M}_i$ has the generalized cylindrical end, and, as we showed in Subsection 4.3, we may assume that the strongly acyclic compatible systems and admissible partitions of unity on $\hat{M}_i$ satisfy Condition (TI). We may also assume that the deformation $\hat{M}_1$ for M_1 is G-equivariant. In addition we can deform P into $\hat{P}$ together with its connection so that it is translationally invariant on the end. Then one can check that the product $\hat{M} = \hat{P} \times_G \hat{M}_1$ has the generalized cylindrical end. Because of the excision property of the local index, we have that M_i (resp. M) and $\hat{M}_i$ (resp. $\hat{M}$) have the same local index. So hereafter we assume that M_i and M has the generalized cylindrical end and all the data satisfy Condition (TI).

Let $\{\rho_{i,\alpha}^2\}_{\alpha\in\widetilde{A}_i}$ be admissible partitions of unity of M_i. We may assume that $\{\rho_{1,\alpha}^2\}$ is G-invariant. Using these partitions of unity we have an admissible partition of unity $\{\rho_{\alpha_0,\alpha_1}^2\}_{(\alpha_0,\alpha_1)\in\widetilde{A}}$ on $M=\cup_{(\alpha_0,\alpha_1)}V_{\alpha_0,\alpha_1}$ which is defined by $\rho_{\alpha_0,\alpha_1}([u,y]) := \rho_{0,\alpha_0}(\pi(u))\rho_{1,\alpha_1}(y)$ for $[u,y]\in M$.

For any translationally invariant Dirac-type operators D_i on $\Gamma(W_i)$, using a local trivialization of P we have their lifts $\widetilde{D}_0\otimes \mathrm{id}_{\widetilde{W}_1}$ and $\epsilon_{\widetilde{W}_0}\otimes\widetilde{D}_1$ on $\Gamma(W)$ as in Subsection 3.6. Note that $D := \widetilde{D}_0\otimes \mathrm{id}_{\widetilde{W}_1} + \epsilon_{\widetilde{W}_0}\otimes\widetilde{D}_1$ is a translationally invariant Dirac-type operator on $\Gamma(W)$.

Because of Lemma 5.5 if we take a positive number t large enough, then the inequality in Proposition 4.10 holds for deformed operators $D_{i,t}$ and D_t on M_i and M. On the other hand we have a decomposition $D_t = D_t^B + D_t^F$, where

$$\begin{aligned}
D_t^B &:= \left(\widetilde{D}_0 + t\sum_{\alpha_0}\left(\sum_{\alpha_1}\rho_{\alpha_1}^2\right)\pi^*\rho_{0,\alpha_0}\widetilde{D}_{0,\alpha_0}\pi^*\rho_{0,\alpha_0}\right)\otimes \mathrm{id}_{\widetilde{W}_1}\\
&= \left(\widetilde{D}_0 + t\sum_{\alpha_0}\pi^*\rho_{0,\alpha_0}\widetilde{D}_{0,\alpha_0}\pi^*\rho_{0,\alpha_0}\right)\otimes \mathrm{id}_{\widetilde{W}_1}\\
D_t^F &:= \epsilon_{\widetilde{W}_0}\otimes\left(\widetilde{D}_1 + t\sum_{\alpha_1}\left(\sum_{\alpha_0}\pi^*\rho_{\alpha_0}^2\right)\rho_{1,\alpha_1}\widetilde{D}_{1,\alpha_1}\rho_{1,\alpha_1}\right)\\
&= \epsilon_{\widetilde{W}_0}\otimes\left(\widetilde{D}_1 + t\sum_{\alpha_1}\rho_{1,\alpha_1}\widetilde{D}_{1,\alpha_1}\rho_{1,\alpha_1}\right).
\end{aligned}$$

Note that D_t^F is a differential operator along fibers of $\pi : M\to M_0$. They anti-commutes each other. Namely,

Lemma 5.6. $D_t^B D_t^F + D_t^F D_t^B = 0$.

Moreover since M_1 and M have the generalized cylindrical end structure, we have the following by Lemma 4.2.

Lemma 5.7. *D_t and D_t^F satisfy Assumption 3.26.*

When we write $\ker D_{1,t} = E^0\oplus E^1$ as the G-equivariant $\mathbb{Z}/2$-graded vector space, G-equivariant local index of (M_1,V_1,W_1) can be written as $\mathrm{ind}_G(M_1,V_1,W_1) = [E^0]-[E^1]\in R(G)$. Let $\underline{E}^i$ be the vector bundle over M_0 defined by $\underline{E}^i = P\times_G E^i$. Then the strongly acyclic compatible system on (M_0,V_0,W_0) induces another strongly acyclic compatible systems on $(M_0, V_0, W_0 \otimes \underline{E}^i)$ via $\{D_{0,\alpha} \otimes \mathrm{id}_{E^i}\}$ for $i = 0, 1$. Lemma 5.6, Lemma 5.7 and the product formula in Section 3 imply the following product formula of local indices.

Theorem 5.8. *We have the following product formula.*

$$\mathrm{ind}(M_0, V_0, W_0 \otimes \underline{E}^0) - \mathrm{ind}(M_0, V_0, W_0 \otimes \underline{E}^1) = \mathrm{ind}(M, V, W) \in \mathbb{Z}.$$

6. Four-dimensional case

6.1. **Local indices for elliptic singularities.** A critical point of a $2n$-dimensional singular Lagrangian fibration $\mu\colon (M, \omega) \to B$ is called a *nondegenerate elliptic singular point of rank* k $(\leq n)$ if there exists a symplectic coordinates $x_1, \dots, x_n, y_1, \dots, y_n$ such that in these coordinate, μ is written as $\mu = (x_1, \dots, x_k, x_{k+1}^2 + y_{k+1}^2, \dots, x_n^2 + y_n^2)$. See [12, 11, 9]. In this subsection we calculate local indices for elliptic singularities in four-dimensional case.

6.1.1. *Definition of* $RR_0(a_1, a_2)$. Let $D := \{z \in \mathbb{C} \mid |z| < 1\}$ be the unit open disc in $\mathbb{C}$. Let X_0 be the product of two copies of D with symplectic structure

$$\omega_0 := \frac{\sqrt{-1}}{2\pi} \sum_{k=1}^{2} dz_k \wedge d\overline{z}_k,$$

and (L_0, ∇^{L_0}) a prequantizing line bundle on (X_0, ω_0).

Let us consider the structure of a singular Lagrangian fibration $\mu_0\colon (X_0, \omega_0) \to [0, 1) \times [0, 1)$ on X_0 which is defined by

$$\mu_0(z) := \left(|z_1|^2, |z_2|^2\right).$$

We put the following assumption.

Assumption 6.1. The cohomology groups $H^*\left(\mu_0^{-1}(b); (L_0, \nabla^{L_0})|_{\mu_0^{-1}(b)}\right)$ vanish for all points $b \in [0, 1) \times [0, 1)$ except for $b = (0, 0)$.

Let a_1 and $a_2 \in \mathbb{Z}$ be arbitrary integers. We define a good compatible fibration on $X_0 \smallsetminus \{(0, 0)\}$ consisting of three quotient maps of the torus actions by

$$\pi_0^0\colon V_0^0 := X_0 \cap (\mathbb{C}^* \times \mathbb{C}^*) \to U_0^0 := V_0^0/T^2,$$
$$\pi_1^0\colon V_1^0 := \{(z_1, z_2) \in X_0 \mid |z_1| > |z_2|, |z_1|^2 + a_1|z_2|^2 \notin \mathbb{Z}\} \to U_1^0 := V_1^0/S^1,$$
$$\pi_2^0\colon V_2^0 := \{(z_1, z_2) \in X_0 \mid |z_1| < |z_2|, a_2|z_1|^2 + |z_2|^2 \notin \mathbb{Z}\} \to U_2^0 := V_2^0/S^1,$$

where the T^2-action on V_0^0 is the standard one, the S^1-action on V_1^0 is defined by

$$t(z_1, z_2) := (tz_1, t^{a_1} z_2), \tag{11}$$

and the S^1-action on V_2^0 is defined by

$$t(z_1, z_2) := (t^{a_2} z_1, tz_2). \tag{12}$$

We take and fix an arbitrary Hermitian structure (g_0, J_0) invariant under the standard T^2-action on X_0 and compatible with ω_0. Since g_0 is T^2-invariant g_0 induces a compatible Riemannian metric of this compatible fibration.

Let W_0 be the Hermitian vector bundle on X_0 which is defined by

$$W_0 := \wedge^{\bullet}_{\mathbb{C}}(TX_0)_{\mathbb{C}} \otimes L_0.$$

W_0 is a $\mathbb{Z}_2$-graded Clifford module bundle with respect to the Clifford module structure (6). We take a compatible system $\{D_i\}_{i=0,1,2}$ to be the family of de Rham operators along fibers of π_i^0 $(i=0,1,2)$ which is defined by the same way as in Example 2.28. By Assumption 6.1 and the definition of V_i^0 $(i=1,2)$ the kernel of all D_i vanish. We should notice that for $z \in V_i^0$ the cohomology groups $H^*\left(\left(\pi_i^0\right)^{-1}\left(\pi_i^0(z)\right); \left(L_0, \nabla^{L_0}\right)|_{\left(\pi_i^0\right)^{-1}\left(\pi_i^0(z)\right)}\right)$ vanish if and only if z satisfies the equality $|z_1|^2 + a_1|z_2|^2 \notin \mathbb{Z}$ for $i=1$ and $a_2|z_1|^2 + |z_2|^2 \notin \mathbb{Z}$ for $i=2$. Hence $\{D_i\}$ is strongly acyclic.

Definition 6.2. Let D be a Dirac-type operator on W_0. We define $RR_0(a_1, a_2)$ to be the local index in the sense of Definition 4.18 with respect to D and the above data.

Remark 6.3. $RR_0(a_1, a_2)$ does not depend on the choice of a compatible Hermitian structure (g_0, J_0) since it is deformation invariant.

6.1.2. *Definition of* $RR_1(a_+, a_-)$. Let $X_1 := (0,1) \times S^1 \times D$ be the product of $(0,1) \times S^1$ and D with symplectic structure

$$\omega_1 := dr \wedge d\theta + \frac{\sqrt{-1}}{2\pi} dz \wedge d\bar{z}$$

for $(r, e^{2\pi\sqrt{-1}\theta}, z) \in X_1$, and (L_1, ∇^{L_1}) a prequantizing line bundle on (X_1, ω_1).

Let us consider the structure of singular Lagrangian fibration $\mu_1 \colon (X_1, \omega_1) \to (0,1) \times [0,1)$ which is defined by

$$\mu_1(r, u, z) := \left(r, |z|^2\right).$$

We put the following assumption.

Assumption 6.4. For all points $b \in (0,1) \times [0,1)$, $H^*(\mu_1^{-1}(b); (L_1, \nabla^{L_1})|_{\mu_1^{-1}(b)})$ vanish.

Let a_+ and $a_- \in \mathbb{Z}$ be arbitrary integers. We take an element $r_1 \in (0,1)$ and fix it. Then, we define a good compatible fibration on $X_1 \smallsetminus \mu_1^{-1}(r_1, 0)$ consisting of three quotient maps of the torus actions by

$$\pi_0^1 \colon V_0^1 := (0,1) \times S^1 \times (D \smallsetminus \{0\}) \to U_0^1 := V_0^1 / T^2,$$
$$\pi_1^1 \colon V_1^1 := \{(r,u,z) \in (r_1, 1) \times S^1 \times D \mid r + a_+|z|^2 \notin \mathbb{Z}\} \to U_1^1 := V_1^1 / S^1,$$
$$\pi_2^1 \colon V_2^1 := \{(r,u,z) \in (0, r_1) \times S^1 \times D \mid r + a_-|z|^2 \notin \mathbb{Z}\} \to U_2^1 := V_2^1 / S^1,$$

where the T^2-action on V_0^1 is defined by

$$t(r,u,z) := (r, t_1 u, t_2 z),$$

the S^1-action on V_1^1 is defined by

$$t(r,u,z) := (r, tu, t^{a_+} z),$$

and the S^1-action on V_2^1 is defined by

$$t(r,u,z) := (r, tu, t^{a_-} z).$$

We take an arbitrary Hermitian structure (g_1, J_1) which is invariant under the standard T^2-action on X_1 and compatible with ω_1 and fix it. We define the $\mathbb{Z}_2$-graded Clifford module bundle W_1 and the strongly acyclic compatible system in the same way as in Section 6.1.1.

Definition 6.5. Let D be a Dirac-type operator on W_1. Then, we define $RR_1(a_+, a_-)$ to be the local index in the sense of Definition 4.18 with respect to D and the above data.

Remark 6.6. $RR_1(a_+, a_-)$ does not depend on the choice of a compatible Hermitian structure (g_1, J_1) since it is deformation invariant.

6.1.3. *Computation.* First we can show the following lemma.

Lemma 6.7. *For integers $a, b, c \in \mathbb{Z}$ we have*

$$RR_0(a, b) = RR_0(b, a), \quad RR_1(a, b) = RR_1(a + c, b + c).$$

Proof. We prove the latter equation. The proof of the former equation is similar. Let $\varphi\colon (0, 1) \times S^1 \times D \to (0, 1) \times S^1 \times D$ be the diffeomorphism which is defined by

$$\varphi(r, u, z) = (r, u, u^c z).$$

On the target space of φ we consider the same compatible fibration as $\{\pi_i^1\}_{i=0,1,2}$ except that a and b are replaced by $a+c$ and $b+c$, respectively. Then φ induces an isomorphism between compatible fibrations.

As the other data on the target space of φ we consider the data which are induced from those on the source space by φ^{-1}. Then the local index for the induced data on the target space is nothing but $RR_1(a, b)$.

On the other hand, the data $(\varphi^{-1})^*\omega_1$ and $(\varphi^{-1})^*\nabla^{L_1}$ can be deformed to ω_1 and ∇^{L_1} by linear deformations. Since the local index is invariant under continuous deformation this implies that the latter equation. □

Moreover, we can also show the following lemma by Theorem 4.19.

Lemma 6.8.

$$RR_0(a, b) = RR_0(a', b) + RR_1(a', a), \qquad RR_1(a, c) = RR_1(a, b) + RR_1(b, c).$$

Then we can calculate $RR_0(a_1, a_2)$ and $RR_1(a_+, a_-)$.

Theorem 6.9.

$$RR_0(a_1, a_2) = 1, \qquad RR_1(a_+, a_-) = 0.$$

Proof. We show $RR_0(0, 1) = 1$ and $RR_0(0, 0) = 1$. Then the theorem follows from these equalities and Lemma 6.7 and 6.8.

First we show $RR_0(0, 1) = 1$. Let us consider the standard toric action on $\mathbb{C}\mathrm{P}^2$ with hyperplane bundle as a prequantizing line bundle. We adopt the moment map μ of this action as a singular Lagrangian fibration. The image B of μ is the triangle in $\mathbb{R}^2$ with vertices $(0, 0)$, $(1, 0)$, $(0, 1)$, and μ has three Bohr-Sommerfeld fibers which corresponds one-to-one to three fixed points $[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]$ of the toric action.

We construct a compatible fibration on $\mathbb{C}P^2 \smallsetminus \{[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]\}$. For each $k \in \mathbb{Z}/3$ let V_k be a pairwise disjoint T^2-invariant open neighborhood of $\{[z_0 : z_1 : z_2] \in \mathbb{C}P^2 \mid z_k = 0\} \smallsetminus \{[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]\}$, and G_k the stabilizer of $\{[z_0 : z_1 : z_2] \in \mathbb{C}P^2 \mid z_k = 0\} \smallsetminus \{[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]\}$. Each G_k is a circle subgroup in T^2 and G_{k-1} acts on V_k freely. Then we put $U_k := V_k/G_{k-1}$ and define $\pi_k\colon V_k \to U_k$ to be the quotient map. We also put $V_4 := U_4 := B \smallsetminus \partial B$ and define $\pi_4\colon V_4 \to U_4$ to be the identity map. These data define a good compatible fibration on $\mathbb{C}P^2 \smallsetminus \{[1 : 0 : 0], [0 : 1 : 0], [0 : 0 : 1]\}$.

The $\mathbb{Z}_2$-graded Clifford module bundle and the strongly acyclic compatible system are defined by the same way as in Section 6.1.1.

Then by Theorem 4.21 the Riemann-Roch number is localized at $[1:0:0], [0:1:0], [0:0:1]$, and the contribution of each fixed point is equal to $RR_0(0,1)$.

On the other hand it is well-known that the Riemann-Roch number of $\mathbb{C}P^2$ is 3. Thus we obtain $RR_0(0,1) = 1$.

Next we show $RR_0(0,0) = 1$. It is a direct consequence of the product formula 5.8 and the fact $[D^+] = 1$(see [4, Theorem 6.7]).

We can also show $RR_0(0,0) = 1$ in the following way. We consider $\mathbb{C}P^1 \times \mathbb{C}P^1$ with standard toric action. The image of the moment map is a square. By the similar construction as above the Riemann-Roch number is localized at four vertices and the contribution of any vertex is $RR_0(0,0)$. On the other hand the Riemann-Roch number of $\mathbb{C}P^1 \times \mathbb{C}P^1$ is four. This implies $RR_0(0,0) = 1$. □

6.2. **Application to locally toric Lagrangian fibrations.** In this subsection we apply the localization formula (Theorem 4.21), the product formula (Theorem 5.8), and Theorem 6.9 to show that for a four-dimensional closed locally toric Lagrangian fibration the Riemann-Roch number is equal to the number of Bohr-Sommerfeld fibers (Theorem 6.23).

6.2.1. *Locally toric Lagrangian fibrations.* Let $\omega_{\mathbb{C}^n}$ be the standard symplectic structure on $\mathbb{C}^n$

$$\omega_{\mathbb{C}^n} := \frac{\sqrt{-1}}{2\pi} \sum_{k=1}^{n} dz_k \wedge d\overline{z}_k.$$

The standard action of T^n on $\mathbb{C}^n$ preserves $\omega_{\mathbb{C}^n}$ and the map $\mu_{\mathbb{C}^n} \colon \mathbb{C}^n \to \mathbb{R}^n$ which is defined by

$$\mu_{\mathbb{C}^n}(z) := \left(|z_1|^2, \dots, |z_n|^2\right)$$

for $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ is a moment map of the standard T^n-action. Note that the image of $\mu_{\mathbb{C}^n}$ is the n-dimensional standard positive cone

$$\mathbb{R}^n_+ := \{r = (r_1, \dots, r_n) \in \mathbb{R}^n \colon r_i \geq 0 \ i = 1, \dots, n\}.$$

Let (M, ω) be a $2n$-dimensional symplectic manifold and B an n-dimensional manifold with corners.

Definition 6.10 ([9, 13])**.** A map $\mu \colon (M, \omega) \to B$ is called a *locally toric Lagrangian fibration* if there exists a system $\{(U_\alpha, \varphi^B_\alpha)\}$ of coordinate neighborhoods of B modeled on $\mathbb{R}^n_+$, and for each α there exists a symplectomorphism $\varphi^M_\alpha \colon (\mu^{-1}(U_\alpha), \omega) \to (\mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha)), \omega_{\mathbb{C}^n})$ such that $\mu_{\mathbb{C}^n} \circ \varphi^M_\alpha = \varphi^B_\alpha \circ \mu$.

Note that a locally toric Lagrangian fibration is a singular Lagrangian fibration that allows only elliptic singularities.

By the definition of a manifold with corners, B is equipped with a natural stratification. We denote by $\mathcal{S}^{(k)}B$ the k-dimensional part of B, namely, $\mathcal{S}^{(k)}B$ consists of those points which have exactly k nonzero components in a local coordinate system. Then, it is easy to see that the fiber of μ at a point in $\mathcal{S}^{(k)}B$ is a k-dimensional torus. In particular, all fibers of μ are smooth.

Example 6.11 (Projective toric variety)**.** The moment map of a nonsingular projective toric variety is a locally toric Lagrangian fibration.

Example 6.12 (Non toric example)**.** Let $c \in \mathbb{N}$ be a positive integer. We consider the diagonal Hamiltonian S^1-action on $(\mathbb{C}^2, \omega_{\mathbb{C}^2})$ with moment map

$$\Phi(z) := \|z\|^2 - c.$$

It is well-known that the symplectic quotient $\left(\Phi^{-1}(0), \omega_{\mathbb{C}^2}|_{\Phi^{-1}(0)}\right)/S^1$ is $\mathbb{C}P^1$ with c times Fubini-Study form ω_{FS}. In the rest of this example we identify $(\mathbb{C}P^1, c\omega_{FS})$ with $\left(\Phi^{-1}(0), \omega_{\mathbb{C}^2}|_{\Phi^{-1}(0)}\right)/S^1$.

Let $\widetilde{\mu}\colon (\widetilde{M}, \widetilde{\omega}) \to \widetilde{B}$ be the singular Lagrangian fibration which is defined by

$$\begin{aligned}
(\widetilde{M}, \widetilde{\omega}) &:= (\mathbb{R} \times S^1 \times \mathbb{C}P^1, dr \wedge d\theta \oplus c\omega_{FS}),\\
\widetilde{B} &:= \mathbb{R} \times [0, c],\\
\widetilde{\mu}(r, u, [z_0 : z_1]) &:= (r, |z_1|^2),
\end{aligned}$$

where we use the coordinate $(r, e^{2\pi\sqrt{-1}\theta}) \in \mathbb{R} \times S^1$. For a negative integer $a \in \mathbb{Z}$ $(a < 0)$ and a positive integer $b \in \mathbb{N}$, we define the $\mathbb{Z}$-actions on $\widetilde{M}$ and $\widetilde{B}$ by

$$n(r, u, [z_0 : z_1]) := \left(r + n(-a|z_1|^2 + b), u, [z_0 : u^{na} z_1]\right), \tag{13}$$

$$n(r_1, r_2) := (r_1 + n(-ar_2 + b), r_2). \tag{14}$$

It is easy to see that (13) and (14) are free $\mathbb{Z}$-actions and (13) preserves $\widetilde{\omega}$. Then we put

$$\begin{aligned}
(M, \omega) &:= (\widetilde{M}, \widetilde{\omega})/\mathbb{Z},\\
B &:= \widetilde{B}/\mathbb{Z}.
\end{aligned}$$

It is also easy to see that $\widetilde{\mu}$ is equivariant with respect to (13) and (14). Hence $\widetilde{\mu}$ induces the map from M to B which we denote by $\mu\colon (M, \omega) \to B$. By construction, B is a cylinder and μ is a locally toric Lagrangian fibration which has singular fibers on ∂B.

Let $\mu\colon (M^{2n}, \omega) \to B$ be a locally toric Lagrangian fibration. By definition, for each α there is a symplectomorphism $\varphi^M_\alpha\colon \mu^{-1}(U_\alpha) \to \mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha))$, and $\mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha))$ has a T^n-action which is obtained by restricting the standard T^n-action on $\mathbb{C}^n$. Then, it is known by [13, Proposition 3.13] that on each nonempty overlap $U_\alpha \cap U_\beta$ there exists an automorphism $\rho_{\alpha\beta} \in \mathrm{Aut}(T^n)$ of T^n such that $\varphi^M_{\alpha\beta} := \varphi^M_\alpha \circ (\varphi^M_\beta)^{-1}$ is $\rho_{\alpha\beta}$-equivariant, namely,

$$\varphi^M_{\alpha\beta}(tx) = \rho_{\alpha\beta}(t)\varphi^M_{\alpha\beta}(x)$$

for $t \in T^2$ and $x \in \mu^{-1}_{\mathbb{C}^n}(\varphi^B_\beta(U_\alpha \cap U_\beta))$. Moreover,we can show that $\rho_{\alpha\beta}$'s form a Čech one-cocycle $\{\rho_{\alpha\beta}\}$ on $\{U_\alpha\}$ with coefficients in $\mathrm{Aut}(T^n)$. Hence it defines an element $[\{\rho_{\alpha\beta}\}]$ in the Čech cohomology $H^1(B; \mathrm{Aut}(T^n))$. Then we have the following lemma.

Lemma 6.13 ([13])**.** *The Čech cohomology class* $[\{\rho_{\alpha\beta}\}]$ *is the obstruction class in order that the* T^n*-actions on* $\mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha))$ *for all* α *can be patched together to obtain a global* T^n*-action on* M*.*

For more detail see [13].

Let $q_B\colon \widetilde{B} \to B$ be the universal covering of B. Since the Čech cohomology $H^1(B; \mathrm{Aut}(T^n))$ is identified with the moduli space of representations of the fundamental group $\pi_1(B)$ of B to $\mathrm{Aut}(T^n)$, the fiber product $q_B^* M := \{(\widetilde{b}, x) \in \widetilde{B} \times M \mid q_B(\widetilde{b}) = \mu(x)\}$ admits a T^n-action.

We take a representative $\rho\colon \pi_1(B) \to \mathrm{Aut}(T^n)$ of the equivalence class of representations corresponding to $[\{\rho_{\alpha\beta}\}]$. Then the T^n-action on q_B^*M can be written explicitly. See [14, Lemma 3.1] for the explicit description.

On the other hand, by the construction, $\pi_1(B)$ acts on q_B^*M from the left by the inverse of the deck transformation, and it is shown that the T^n-action and the $\pi_1(B)$-action satisfy the following relationship

$$t(a\widetilde{x}) = a\left(\rho(a^{-1})(t)\widetilde{x}\right) \tag{15}$$

for $t \in T^n$, $a \in \pi_1(B)$, and $\widetilde{x} \in q_B^*M$. Let $T^n \rtimes_\rho \pi_1(B)$ be the semidirect product of T^n and $\pi_1(B)$ with respect to ρ. Then, (15) implies that these actions form an action of $T^n \rtimes_\rho \pi_1(B)$ on q_B^*M. For more details see [14].

Let $q_M\colon q_B^*M \to M$ be the natural projection. Note that $q_M^*\omega$ is $T^n \rtimes_\rho \pi_1(B)$-invariant since ω is invariant under the T^n-action on $\mu^{-1}(U_\alpha)$ induced by the standard T^n-action on $\mathbb{C}^n$ for each α. Now we show the following lemma.

Lemma 6.14. *There exists a Hermitian structure $(\widetilde{g}, \widetilde{J})$ on q_B^*M compatible with $q_M^*\omega$ which is invariant under the action of $T^n \rtimes_\rho \pi_1(B)$.*

Proof. It is sufficient to show that the existence of an invariant Riemannian metric. Let g' be a Riemannian metric on M. We define the Riemannian metric $\widetilde{g}$ on q_B^*M by

$$\widetilde{g}_{\widetilde{x}}(u,v) := \int_{T^n} \left(\varphi_t^*(q_M^*g')\right)_{\widetilde{x}}(u,v)dt,$$

where φ_t implies the T^n-action for $t \in T^n$. It is sufficient to show that $\widetilde{g}$ is $\pi_1(B)$-invariant. For $a \in \pi_1(B)$ we denote the $\pi_1(B)$-action by ϕ_a. Then we have

$$\begin{aligned}
(\phi_a^*\widetilde{g})_{\widetilde{x}}(u,v) &= \int_{T^n} \left(\phi_a^*\left(\varphi_t^*(q_M^*g')\right)\right)_{\widetilde{x}}(u,v)dt \\
&= \int_{T^n} \left(\varphi_{\rho(a^{-1})(t)}^*\left(\phi_a^*(q_M^*g')\right)\right)_{\widetilde{x}}(u,v)dt \\
&= \int_{T^n} \left(\varphi_{\rho(a^{-1})(t)}^*(q_M^*g')\right)_{\widetilde{x}}(u,v)dt \\
&= \det\rho(a^{-1}) \int_{T^n} \left(\varphi_{\rho(a^{-1})(t)}^*(q_M^*g')\right)_{\widetilde{x}}(u,v)\rho(a^{-1})^*dt \\
&= \int_{T^n} \left(\varphi_t^*(q_M^*g')\right)_{\widetilde{x}}(u,v)dt. \\
&= \widetilde{g}_{\widetilde{x}}(u,v).
\end{aligned}$$

Here we remark that $\det\rho(a^{-1}) = \pm 1$ since $\rho(a^{-1}) \in \mathrm{Aut}(T^n)$. ☐

Corollary 6.15 (the existence of an invariant Hermitian structure)**.** *There exists a Hermitian structure (g, J) on M compatible with ω such that on each $\mu^{-1}(U_\alpha)$ (g, J) is invariant under the T^n-action on $\mu^{-1}(U_\alpha)$ which is induced from the T^n-action on $\mu_{\mathbb{C}^n}^{-1}(\varphi_\alpha^B(U_\alpha))$ with the identification φ_α^M.*

Proof. By Lemma 6.14 there is a $T^n \rtimes_\rho \pi_1(B)$-invariant Hermitian structure $(\widetilde{g}, \widetilde{J})$ on q_B^*M compatible with $q_M^*\omega$. In particular, since $(\widetilde{g}, \widetilde{J})$ is $\pi_1(B)$-invariant, $(\widetilde{g}, \widetilde{J})$ induces an ω-compatible Hermitian structure on M which is denoted by (g, J). Then, (g, J) is the required one. ☐

Lemma 6.16 (The existence of an averaging operation)**.** *Suppose that there exists a compatible fibration* $\{\pi_\alpha\colon V_\alpha\to U_\alpha\}$ *on* M *such that for each* α *a fiber of* π_α *is contained in that of* μ*, namely,* $\pi_\alpha^{-1}\pi_\alpha(x)\subset\mu^{-1}\mu(x)$ *for* $x\in V_\alpha$*. There exists an averaging operation* $I\colon C^\infty(M)\to C^\infty(M)$ *with respect to* $\{\pi_\alpha\colon V_\alpha\to U_\alpha\}$*.*

Proof. For $f\in C^\infty(M)$ let $\widetilde{f}\in C^\infty(q_B^*M)$ be the function on q_B^*M which is defined by

$$\widetilde{f}(\widetilde{x}):=\int_{T^n}(f\circ q_M)(t\widetilde{x})dt.$$

Then, by the similar way to that in the proof of Lemma 6.14, we can show that $\widetilde{f}$ is $T^n\rtimes_\rho\pi_1(B)$-invariant. Hence it descends to the function on M. We denote it by $I(f)$. Then, it is clear that $I(f)$ satisfies the properties in Definition 2.11. □

6.2.2. *Bohr-Sommerfeld fibers and the Riemann-Roch number.* Let $\mu\colon(M,\omega)\to B$ be a prequantized locally toric Lagrangian fibration with prequantizing line bundle (L,∇). Recall that, as described above, all fibers are smooth.

Definition 6.17. A fiber F of μ is said to be *Bohr-Sommerfeld* if the restriction $(L,\nabla)|_F$ is trivially flat. A point b of B is also said to be *Bohr-Sommerfeld* if the fiber $\mu^{-1}(b)$ is Bohr-Sommerfeld.

Remark 6.18. A fiber F of μ is Bohr-Sommerfeld if and only if the cohomology $H^*(F;(L,\nabla)|_F)$ does not vanish, see Lemma 2.29. This is also equivalent to the condition that the de Rham operator on F with coefficients in $(L,\nabla)|_F$ has non zero kernel.

First we specify Bohr-Sommerfeld points for the local model.

Proposition 6.19. *Let* (L,∇) *be a prequantizing line bundle on* $(\mathbb{C}^n,\omega_{\mathbb{C}^n})$*. Then, a point* $b\in\mathbb{R}^n_+$ *is Bohr-Sommerfeld if and only if* $b\in\mathbb{R}^n_+\cap\mathbb{Z}^n$*.*

Proof. Since $\mathbb{C}^n$ is contractible L is trivial as a complex line bundle. Then we can assume that L is of the form $L=\mathbb{C}^n\times\mathbb{C}$ without loss of generality. Then, ∇ can be written as

$$\nabla=d-2\pi\sqrt{-1}A$$

for some one form on $\mathbb{C}^n$ with $dA=\omega_{\mathbb{C}^n}$. Moreover A is unique up to exact one form since $\mathbb{C}^n$ is contractible. In particular, A is of the form

$$A=\frac{\sqrt{-1}}{4\pi}\sum_{i=1}^n(z_id\bar{z}_i-\bar{z}_idz_i)+df$$

for some smooth function f on $\mathbb{C}^n$.

By using the polar coordinate $z_i=r_ie^{2\pi\sqrt{-1}\theta_i}$ we can write $\mu_{\mathbb{C}}$ and A in the following forms

$$\mu_{\mathbb{C}^n}=(r_1^2,\dots,r_n^2),\ A=\sum_i r_i^2d\theta_i+df.$$

In particular, we see that the tangent space along a nonsingular fiber of $\mu_{\mathbb{C}^n}$ is spanned by ∂_{θ_i}'s. Thus a direct computation shows that a point $b\in\mathbb{R}^n_+$ is Bohr-Sommerfeld if and only if $b\in\mathbb{R}^n_+\cap\mathbb{Z}^n$. □

By the above proposition and the definition of a locally toric Lagrangian fibration we can obtain the following corollary.

Corollary 6.20. *For a locally toric Lagrangian fibration Bohr-Sommerfeld fibers appear discretely.*

Example 6.21. For a nonsingular projective toric variety it is well-known that Bohr-Sommerfeld fibers correspond one-to-one to the integral points in the moment polytope. For example see [3].

Example 6.22. We consider the locally toric Lagrangian fibration $\mu\colon (M,\omega)\to B$ in Example 6.12. We show that (M,ω) is prequantizable.

Let (H_c,∇^{H_c}) be the c times tensor power of the hyperplane bundle on $\mathbb{C}P^1$. With the identification of $(\mathbb{C}P^1, c\omega_{FS})$ and the symplectic quotient $\left(\Phi^{-1}(0),\omega_{\mathbb{C}^2}|_{\Phi^{-1}(0)}\right)/S^1$ in Example 6.12 (H_c,∇^{H_c}) can be written in the following explicit way

$$(H_c,\nabla^{H_c}) = \left(\Phi^{-1}(0)\times\mathbb{C}, d+1/2\sum_i(z_i d\overline{z}_i - \overline{z}_i dz_i)\right)/S^1,$$

where the S^1-action is defined by

$$t\cdot(z_0,z_1,w) := (tz_0, tz_1, t^c w).$$

Now let us define the prequantizing line bundle $(\widetilde{L},\widetilde{\nabla})$ on $(\widetilde{M},\widetilde{\omega})$ by

$$(\widetilde{L},\widetilde{\nabla}) := \left(\mathrm{pr}_1^*(\mathbb{R}\times S^1\times\mathbb{C}, d-2\pi\sqrt{-1}rd\theta)\otimes_{\mathbb{C}}\mathrm{pr}_2^*(H_c,\nabla^{H_c})\right).$$

We also define the lift of the $\mathbb{Z}$-action (13) on $\widetilde{M}$ to $\widetilde{L}$ by

$$n(r,u,[z_0:z_1,w]) := \left(r+n(-a|z_1|^2+b), u, [z_0:u^{na}z_1, u^{nb}w]\right). \tag{16}$$

It is easy to see that (16) preserves $\widetilde{\nabla}$ and the standard Hermitian metric. We put

$$(L,\nabla) := (\widetilde{L},\widetilde{\nabla})/\mathbb{Z}.$$

Then (L,∇) is a prequantizing line bundle on (M,ω).

Next we see the Bohr-Sommerfeld fibers of μ with respect to (L,∇). The direct computation shows that Bohr-Sommerfeld fibers of $\widetilde{\mu}$ correspond one-to-one to the elements in $\widetilde{B}\cap\mathbb{Z}^2$. Let F be a fundamental domain of the $\mathbb{Z}$-action (14) on $\widetilde{B}$. F is written as

$$F := \{(r_1,r_2)\in\widetilde{B}\mid 0\le r_2\le c,\ -1/2\le r_1 < -ar_2+b-1/2\}.$$

Then, Bohr-Sommerfeld fibers of μ correspond one-to-one to the elements in $F\cap\mathbb{Z}^2$. See Figure 1.

In the rest of this section we assume that M is closed. Let (g,J) be a Hermitian structure on M compatible with ω as in Corollary 6.15. We define the Hermitian vector bundle W on M by

$$W := \wedge^\bullet_{\mathbb{C}}TM_{\mathbb{C}}\otimes L. \tag{17}$$

W is a $\mathbb{Z}_2$-graded Clifford module bundle with respect to the Clifford module structure (6). Let D be the Dirac-type operator on W. We define the *Riemann-Roch number* to be the index of D.

The purpose of this section is to show the following theorem.

Theorem 6.23. *Let $\mu\colon (M,\omega)\to B$ be a four-dimensional prequantizable locally toric Lagrangian fibration with prequantizing line bundle (L,∇). Then the Riemann-Roch number is equal to the number of both nonsingular and singular Bohr-Sommerfeld fibers.*



Figure 1. Bohr-Sommerfeld points in Example 6.12

Proof. Let B_{BS} be the set of Bohr-Sommerfeld points of μ in B. We put $V := \mu^{-1}(B \smallsetminus B_{BS})$. In order to prove Theorem 6.23 we define a good compatible fibration on V as follows.

On the regular non Bohr-Sommerfeld points $U_0 := \mathcal{S}^{(2)}B \smallsetminus B_{BS}$ of μ we define the fibration by

$$\pi_0 := \mu|_{V_0} \colon V_0 := \mu^{-1}(U_0) \to U_0.$$

Since B is compact, there are only finitely many Bohr-Sommerfeld points in $\mathcal{S}^{(1)}B$. Suppose we have exactly k Bohr-Sommerfeld points $p_1, \dots, p_k$ in $\mathcal{S}^{(1)}B$, namely,

$$\{p_1, \dots, p_k\} = B_{BS} \cap \mathcal{S}^{(1)}B.$$

For each i we take a contractible open neighborhood W_i of p_i in B which satisfies the following properties.

1. For each i W_i contains no Bohr-Sommerfeld points except for p_i.
2. For each i $W_i \cap \mathcal{S}^{(1)}B$ is connected.
3. For each i W_i does not intersect $\mathcal{S}^{(0)}B$, namely, $W_i \cap \mathcal{S}^{(0)}B = \emptyset$.
4. W_i's are pairwise disjoint, namely, $W_i \cap W_j = \emptyset$ for all $i \neq j$.
5. There exist finitely many non Bohr-Sommerfeld points in $\mathcal{S}^{(1)}B$, say $q_1, \dots, q_l$, such that we have
$$\bigcup_{i=1}^{k} W_i \cap \mathcal{S}^{(1)}B = \mathcal{S}^{(1)}B \smallsetminus \{q_1, \dots, q_l\}.$$

It is possible to take such neighborhoods since a connected component of ∂B is compact.

We put $V_i' := \mu^{-1}(W_i)$. Since W_i is contractible, by [13, Proposition 3.5], there exists a T^2-action on V_i'. Moreover, there exist a coordinate neighborhood $(U_{\alpha_i}, \varphi^B_{\alpha_i})$ of B containing p_i, a diffeomorphism $\varphi^M_{\alpha_i} \colon \mu^{-1}(U_{\alpha_i}) \to \mu^{-1}_{\mathbb{C}^2}(\varphi^B_{\alpha_i}(U_{\alpha_i}))$ in Definition 6.10, and an automorphism $\rho_{\alpha_i} \in \mathrm{Aut}(T^2)$ which satisfy the following properties.

- $\mu_{\mathbb{C}^2} \circ \varphi^M_{\alpha_i} = \varphi^B_{\alpha_i} \circ \mu$.
- On $V_i' \cap \mu^{-1}(U_{\alpha_i})$ $\varphi^M_{\alpha_i}$ is ρ_{α_i}-equivariant with respect to the T^2-action on V_i' and the standard T^2-action on $\mathbb{C}^2$.



FIGURE 2. W_i's in Example 6.12

Let $\varphi^B_{\alpha_i}(p_i) = (r_1, r_2) \in \mathbb{R}^2_+$. Since $p_i \in \mathcal{S}^{(1)}B$ there exists a unique coordinate r_{j_i} such that $r_{j_i} = 0$. We define the circle subgroup T_i of T^2 by

$$T_i := \rho^{-1}_{\alpha_i}\left(\{t = (t_1, t_2) \in T^2 \mid t_{j_i} = e\}\right).$$

By the properties 2 and 3 of W_i, T_i acts on V_i' freely. Then, for each i we define the open set V_i to be the complement of all T_i-orbits in V_i' on each of which (L, ∇) has a non-trivial global parallel section. We also define the fibration $\pi_i \colon V_i \to U_i$ to be the natural projection

$$\pi_i \colon V_i \to U_i := V_i/T_i$$

with respect to the T_i-action on V_i. Note that for each i we have the following equality

$$V_i \cap \mu^{-1}(\mathcal{S}^{(1)}B) = \mu^{-1}(W_i \cap \mathcal{S}^{(1)}B) \smallsetminus \mu^{-1}(p_i).$$

In order to show this equality, it is sufficient to prove the following claim: for a point $x \in \mu^{-1}(W_i \cap \mathcal{S}^{(1)}B)$ the restriction of (L, ∇) to the fiber $\mu^{-1}(\mu(x))$ of μ is trivially flat if and only if the restriction of (L, ∇) to the T_i-orbit $T_i x$ through x is trivially flat. This can be shown as follows. By the property 2 of W_i, the T^2-action on $\mu^{-1}(W_i \cap \mathcal{S}^{(1)}B)$ has the unique one-dimensional stabilizer $S \subset T^2$. For example, in case of $j_i = 2$, S is written as

$$S = \rho^{-1}_{\alpha_i}\left(\{t = (t_1, t_2) \in T^2 \mid t_1 = e\}\right).$$

In any case, by the definition of T_i, T^2 is decomposed as $T^2 = T_i \times S$. Then, the above claim follows from the fact that in the T^2-action on $\mu^{-1}(\mu(x))$ S acts trivially and T_i acts freely on $\mu^{-1}(\mu(x))$.

By construction $\{\pi_i \colon V_i \to U_i \mid i = 0, \dots, k\}$ is a good compatible fibration on V. Moreover, by Lemma 6.16, there is an averaging operation with respect to $\{\pi_i \colon V_i \to U_i \mid i = 0, \dots, k\}$.

Recall that (g, J) is a Hermitian structure on M compatible with ω as in Corollary 6.15. Then, as in the case of usual torus actions, g defines the compatible Riemannian metric of $\{\pi_i \colon V_i \to U_i \mid i = 0, \dots, k\}$ whose restriction to each fiber of μ is flat, and the $\mathbb{Z}_2$-graded Clifford module bundle W defined by (17) becomes a compatible Clifford module bundle in the sense of Definition 2.19. We define the strongly acyclic compatible system in the same way as in Section 6.1.1. Then by Theorem 4.21, the Riemann-Roch number is localized at Bohr-Sommerfeld fibers and the fibers at $q_1, \dots, q_l$.

We consider their contributions. Since a fiber of μ is connected, by Theorem [4, Theorem 6.11], the contribution of a regular Bohr-Sommerfeld fiber is equal to one.

Next we consider the contributions of singular Bohr-Sommerfeld fibers. By Definition each fiber on $\mathcal{S}^{(0)}B$ is Bohr-Sommerfeld, and its contribution is $RR_0(a_1, a_2)$ for some a_1 and a_2. By Theorem 6.9 it is equal to one.

By the construction of the compatible fibration the local Riemann-Roch number for each singular Bohr-Sommerfeld fiber on $\mathcal{S}^{(1)}B$ is obtained from $[BS^+]$ and $[D^+]$ in [4, Theorem 6.7] by the product formula 5.8. It is also one.

Finally it is easy to see that the contribution of each fibers at $q_1, \dots, q_l$ is equal to $RR_1(a_+, a_-)$ in Section 6.1.2 for some a_+ and a_-. Then by Theorem 6.9 it is zero. This proves Theorem 6.23. □

Example 6.24. Theorem 6.23 recovers Danilov's result [3], which says that for a non-singular projective toric variety the Riemann-Roch number is equal to the number of the lattice points in the moment polytope, in the four-dimensional case.

Example 6.25. As we described in Example 6.22 the Bohr-Sommerfeld fibers correspond one-to-one to the elements in $F \cap \mathbb{Z}^2$. Then by Theorem 6.23 the Riemann-Roch number of (M, ω) is equal to the number of the elements in $F \cap \mathbb{Z}^2$ which is $(c+1)(2b-ac)/2$.

Appendix A. Proof of Proposition 2.15

In this appendix we give a proof of Proposition 2.15. We first give the precise statement.

Let $\{\pi_\alpha : V_\alpha \to U_\alpha\}_\alpha$ be a good compatible fibraiton on M. Let $\{V'_\alpha\}$ be an admissible open covering of M in the second condition in Assumption 2.12. For each $\alpha \in A$ take an open subset U''_α of U_α such that $\pi_\alpha(V'_\alpha) \Subset U''_\alpha \Subset U_\alpha$. We put $V''_\alpha := \pi_\alpha^{-1}(U''_\alpha)$. For each α take a smooth function $\tau_\alpha : M \to \mathbb{R}$ such that $\tau_\alpha \equiv 1$ on V'_α and $\tau_\alpha \equiv 0$ on $M \smallsetminus V''_\alpha$. In this appendix we show the following.

Proposition A.1. *If $\{\pi_\alpha\}$ is a good compatible fibration, then there exists an averaging operation $I : C^\infty(M) \to C^\infty(M)$ such that for all $f \in C^\infty(M)$ and $x \in M$ we have*

$$\min_{y \in \pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)} f(y) \le I(f)(x) \le \max_{y \in \pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)} f(y),$$

where $\alpha''_x := \max\{\alpha \in A \mid x \in V''_\alpha\}$.

Proof. For each $\alpha \in A$ we define a linear map $I_\alpha : C^\infty(M) \to C^\infty(M)$ by

$$I_\alpha(f)(x) := (1 - \tau_\alpha(x))f(x) + \tau_\alpha(x)I^0_\alpha(f)(x),$$

where $I^0_\alpha(f)$ is the integration along fibers of $\pi_\alpha : V_\alpha \to U_\alpha$ with the normalization condition $I^0_\alpha(1) \equiv 1$. Fix any total order of $A = \{\alpha_1, \cdots, \alpha_n\}$ so that if $V_{\alpha_i} \cap V_{\alpha_j} \neq \emptyset$ and the dimension of the fiber of π_{α_i} is bigger than that of π_{α_j} then $i > j$. Using this total order we may write $A = \{1, 2, \cdots, n\}$, $\pi_{\alpha_k} = \pi_k$ and so on. We define a linear map $I : C^\infty(M) \to C^\infty(M)$ by

$$I(f) := I_1 I_2 \cdots I_{n-1} I_n(f).$$

Note that I satisfies the conditions (2) and (3) in Definition 2.11. We show that I satisfies the conditions (1), (4), and (5) respectively in the following three lemmas. □

Lemma A.2. *$I(f)$ is an admissible function for all $f \in C^\infty(M)$. More precisely, if $x, y \in V'_k$ and $\pi_k(x) = \pi_k(y)$ for some $k \in A$, then we have $I(f)(x) = I(f)(y)$.*

Proof. Note that, by definition, if $x, y \in V'_k$ and $\pi_k(x) = \pi_k(y)$ for some $k \in A$, then we have $k \le \alpha''_x = \alpha''_y$ and $\pi_\alpha(x) = \pi_\alpha(y)$ for any $\alpha \le \alpha''_x$ and $x, y \in V_\alpha$. If $\alpha > \alpha''_x$ and $x' \in \pi_\alpha^{-1}\pi_\alpha(x)$, then we have $x' \notin V''_\alpha$, and hence, $I_\alpha(g)(x') = g(x')$ for any smooth function g. In particular we have $I_{\alpha''_x} I_\alpha(g)(x) = I_{\alpha''_x}(g)(x)$. Put $\alpha'_x := \max\{\alpha \in A \mid x \in V'_\alpha\}$. Note that $\alpha'_x = \alpha'_y$. If $\beta < \alpha'_x$ and $x \in V_\beta$, then since $I_{\alpha'_x}(g)$ is constant along $\pi_\beta^{-1}\pi_\beta(x) \subset V'_{\alpha'_x}$ we have $I_\beta I_{\alpha'_x}(g)(x) = I_{\alpha'_x}(g)(x)$. By induction we have

$$I(f)(x) = I_{\alpha'_x} \cdots I_\beta \cdots I_{\alpha''_x}(f)(x),$$

where we may assume that each $\beta \in A$ satisfies $\alpha'_x \le \beta \le \alpha''_x$ and $x, y \in V''_\beta$. Since $x, y \in V'_{\alpha_x}$ and $\pi_{\alpha'_x}(x) = \pi_{\alpha'_x}(y)$ we have

$$I(f)(x) = I^0_{\alpha'_x}(\cdots I_\beta \cdots I_{\alpha''_x}(f))(x) = I^0_{\alpha'_y}(\cdots I_\beta \cdots I_{\alpha''_y}(f))(y) = I(f)(y).$$

□

Lemma A.3. *Let $f : M \to \mathbb{R}$ be a smooth function and $x \in M$. Then we have*

$$\min_{y \in \pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)} f(y) \le I(f)(x) \le \max_{y \in \pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)} f(y),$$

where $\alpha''_x := \max\{\alpha \in A \mid x \in V''_\alpha\}$.

Proof. Note that

$$I_\alpha(g)(x) = (1 - \tau_\alpha(x))g(x) + \tau_\alpha(x) I^0_\alpha(g)(x) \le \max_{\pi_\alpha^{-1}\pi_\alpha(x)} g \tag{18}$$

for any smooth function $g : M \to \mathbb{R}$, $\alpha \in A$ and $x \in V_\alpha$. On the other hand as we showed in the proof of Lemma A.2, we have

$$I(f)(x) = I_{\alpha'_x} \cdots I_\beta \cdots I_{\alpha''_x}(f)(x)$$

for any $x \in M$. Since $\pi^{-1}_{\alpha'_x}\pi_{\alpha'_x}(x) \subset \cdots \pi_\beta^{-1}\pi_\beta(x) \subset \cdots \subset \pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)$ we have

$$I(f)(x) \le \max_{\pi^{-1}_{\alpha'_x}\pi_{\alpha'_x}(x)} \big(\cdots I_\beta \cdots I_{\alpha''_x}(f)\big) \le \cdots \le \max_{\pi^{-1}_{\alpha''_x}\pi_{\alpha''_x}(x)} f$$

by (18) and induction. □

Lemma A.4. *Let $f : M \to \mathbb{R}$ be a smooth function. If $supp f$ is contained in V'_α for some $\alpha \in A$, then $supp I(f)$ is also contained in V'_α.*

Proof. Let $f : M \to \mathbb{R}$ be a smooth function such that $supp f \subset V'_\alpha$ for some $\alpha \in A$. It is enough to show that $supp I_\beta(f) \subset V'_\alpha$ for each $\beta \in A$. Take $x \in supp I_\beta(f)$ and a sequence $\{x_n\}$ in M which satisfies $I_\beta(f)(x_n) \ne 0$ and converges to x. By definition of $I_\beta(f)$ we have $f(x_n) \ne 0$ for infinitely many n or $\tau_\beta(x_n) I^0_\beta(f)(x_n) \ne 0$ for infinitely many n. The former case implies that $x \in supp f \subset V'_\alpha$. In the latter case by taking a subsequence we may assume that $x_n \in V''_\beta$ and $I^0_\beta(f)(x_n) \ne 0$ for all n. In particular we have $x \in \overline{V''_\beta} \subset V_\beta$. Since I^0_β is the integration along fibers there exist a sequence $\{y_n \in \pi_\beta^{-1}\pi_\beta(x_n)\}$ such that $f(y_n) \ne 0$ for all n. By taking a subsequence we may assume $\{y_n\}$ converges to some $y \in \pi_\beta^{-1}\pi_\beta(x) \cap supp f \subset V'_\alpha$. Since $\pi_\beta^{-1}\pi_\beta(V'_\alpha \cap V_\beta) = V'_\alpha \cap V_\beta$ we have $x \in \pi_\beta^{-1}\pi_\beta(x) \subset V'_\alpha$. □

Appendix B. A sufficient condition for Assumption 2.12 (2)

We give a sufficient condition for Assumption 2.12 (2).

Lemma B.1. *Let* $\{\pi_\alpha\}_{\alpha\in A}$ *be a compatible fibration on* M *such that* A *is a finite set and the condition 5' in Definition 2.14 is satisfied. If each* π_α *has a continuous extension as a fiber bundle to the closure of* V_α *with the condition*

$$V_\alpha \cap \overline{V_\beta} = \pi_\beta^{-1}\pi_\beta(V_\alpha \cap \overline{V_\beta}) \tag{19}$$

for all $\beta \in A$*, then* $\{\pi_\alpha\}$ *satisfies the second condition in Assumption 2.12. Namely there exist an admissible open covering* $\{V'_\alpha \mid \alpha \in A\}$ *of* M *such that* $\overline{V}'_\alpha \subset V_\alpha$*.*

Proof. Take and fix any open covering $\{W_\alpha\}_{\alpha\in A}$ of M which satisfies $\overline{W}_\alpha \subset V_\alpha$. Fix any total order of $A = \{\alpha_1, \cdots, \alpha_n\}$ so that if $V_{\alpha_i} \cap V_{\alpha_j} \neq \emptyset$ and the dimension of π_{α_i} is bigger than that of π_{α_j} then $i > j$. Fix $\alpha \in A$ and we define an increasing sequence of open sets $V_\alpha^{(k)} \subset V_\alpha$ inductively in the following way:

$$\begin{aligned}
V_\alpha^{(0)} &:= W_\alpha \\
&\ \vdots \\
V_\alpha^{(k)} &:= \pi_{\alpha_k}^{-1}\pi_{\alpha_k}(V_\alpha^{(k-1)} \cap V_{\alpha_k}) \cup V_\alpha^{(k-1)} \\
&\ \vdots \\
V_\alpha^{(n)} &:= \pi_{\alpha_n}^{-1}\pi_{\alpha_n}(V_\alpha^{(n-1)} \cap V_{\alpha_n}) \cup V_\alpha^{(n-1)}.
\end{aligned}$$

By the construction $\{V'_\alpha := V_\alpha^{(n)}\}_{\alpha\in A}$ is an admissible open covering of M. We show $\overline{V}_\alpha^{(k)} \subset V_\alpha$ by induction on k. Suppose that $\{p_i\}_{i\in\mathbb{N}}$ is a sequence in $V_\alpha^{(k)}$ which converges to p_∞ in M. It is enough to show that if $p_i \in \pi_{\alpha_k}^{-1}\pi_{\alpha_k}(V_\alpha^{(k-1)} \cap V_{\alpha_k})$ for all i then $p_\infty \in V_\alpha$. In this case we have $p_\infty \in \overline{\pi_{\alpha_k}^{-1}\pi_{\alpha_k}(V_\alpha^{(k-1)} \cap V_{\alpha_k})}$. On the other hand since the fibers are compact, π_{α_k} is a closed map. Using the assumption (19) we have

$$\overline{\pi_{\alpha_k}^{-1}\pi_{\alpha_k}(V_\alpha^{(k-1)} \cap V_{\alpha_k})} \subset \pi_{\alpha_k}^{-1}(\overline{\pi_{\alpha_k}(V_\alpha^{(k-1)} \cap V_{\alpha_k})}) \subset \pi_{\alpha_k}^{-1}\pi_{\alpha_k}(\overline{V_\alpha^{(k-1)}} \cap \overline{V_{\alpha_k}}) \subset V_\alpha \cap \overline{V_{\alpha_k}}.$$

In particular we have $p_\infty \in V_\alpha$. □

Remark B.2. Since $\overline{V'_\alpha} \subset V_\alpha$, one can check that $\{V'_\alpha\}_\alpha$ satisfies the same condition as in Assumption 19, i.e., $V'_\alpha \cap \overline{V'_\beta} = \pi_\beta^{-1}\pi_\beta(V'_\alpha \cap \overline{V'_\beta})$ for all $\alpha, \beta \in A$.

Appendix C. Proof of Lemma 2.30

Proof of Lemma 2.30. Let $H_1, H_2, \ldots, H_m$ be the elements of A. Without loss of generality we assume that $H_i \supset H_j$ implies $i \leq j$. We construct a family of open sets $V_i^{(j)}$ $(1 \leq i \leq j \leq m)$ by induction on $1 \leq i \leq m$. For the construction with $i = i_0$ we assume the following properties.

(C1) $V_i^{(j)}$ contains the closure of $V_i^{(j+1)}$ for all $1 \leq i < i_0$ and $i \leq j < m$.
(C2) If $x \in V_i^{(i)}$ for $1 \leq i < i_0$, then we have $G_x \subset H_i$.
(C3) For $x \in M$ with $G_x = H_i$ for some $1 \leq i < i_0$, we have

$$x \in \bigcup_{\{j \mid H_j \supset H_i\}} V_j^{(m)}.$$

(C4) If the intersection $V_i^{(j)} \cap V_j^{(j)}$ is not empty for $1 \leq i < j < i_0$, then we have $H_i \supset H_j$.

If $i_0 = 1$, then the above is the empty assumption. For $1 \leq i_0 \leq m$, using the above properties as the assumption of induction, we construct $V_{i_0}^{(j)}$ $(i_0 \leq j \leq m)$ which satisfy the above properties with replacement of i_0 by $i_0 + 1$.

Suppose $1 \leq i_0 < m$ and assume (C1),(C2),(C3) and (C4). Then (C3) implies that the closed set

$$K_{i_0} := M^{H_{i_0}} \setminus \bigcup_{\{k \,|\, H_k \supsetneq H_{i_0}\}} V_k^{(m)}$$

is contained in $\{x \in M \,|\, G_x = H_{i_0}\}$, where $M^{H_{i_0}}$ is the fixed point set $M^{H_{i_0}} = \{x \in M \,|\, G_x \supset H_{i_0}\}$. Hence (C2) implies that K_{i_0} does not intersect with the open set

$$\bigcup_{\{j < i_0 \,|\, H_{i_0} \not\subset H_j\}} V_j^{(i_0 - 1)}.$$

Let L_{i_0} be the closure of

$$\bigcup_{\{j < i_0 \,|\, H_{i_0} \not\subset H_j\}} V_j^{(i_0)}.$$

Then (C1) implies $K_{i_0} \cap L_{i_0} = \emptyset$. Since K_{i_0} is a subset of $\{x \in M \,|\, G_x = H_{i_0}\}$, there is an open neighborhood V of the closed set K_{i_0} in the complement of L_{i_0} such that for each $x \in V$ we have $G_x \subset H_{i_0}$. Now we take a decreasing sequence of open neighborhoods $V_{i_0}^{(j)}$ $(i_0 \leq j \leq m)$ of K_{i_0} so that $V_{i_0}^{(i_0)} = V$, $V_{i_0}^{(m)} \supset K_{i_0}$ and $V_{i_0}^{(j)}$ contains the closure of $V_{i_0}^{(j+1)}$ for $i_0 \leq j < m$. We can choose the decreasing sequence so that the open sets $V_{i_0}^{(j)}$ $(i_0 \leq j \leq m)$ are G-invariant because the quotient space M/G is a regular space.

Then it is straightforward to check (C1),(C2),(C3) and (C4) are satisfied with i_0 replaced by $i_0 + 1$.

The family of open sets $\{V_{H_i} := V_i^{(m)}\}_{1 \leq i \leq m}$ is an open covering of M and satisfies the required properties.

Now if a family of open sets $\{V_x\}_{x \in M}$ satisfying $gV_x = V_{gx}$ for all $g \in G$ is given, then since K_{i_0} is contained in $\displaystyle\bigcup_{\{x \,\mid\, G_x = H_{i_0}\}} V_x$ we can take a G-invariant open neighborhood $V = V_{i_0}^{(i_0)}$ of K_{i_0} so that $V \subset \bigcup_{G_x = H_{i_0}} V_x$, and hence, we have the required open covering $\{V_H\}_{H \in A}$. □

Appendix D. Proof of Lemma 3.10

Proof of Lemma 3.10. If there is a function f satisfying the property in (1), then $\rho_{\epsilon,a}$ is constructed as follows: For each $\epsilon > 0$ let $\rho_\epsilon : \mathbb{R} \to [0,1]$ be a smooth non-increasing function such that $\rho_\epsilon(l) = 1$ for $l \leq 0$, $\rho_\epsilon(l) = 0$ for $l \geq 2/\epsilon$ and $|d\rho_\epsilon(l)| < \epsilon$ for $i \in \mathbb{R}$. Then the composition $\widetilde{\rho}_{a,\epsilon}(x) = \rho_\epsilon(f(x) - a)$ has the required properties.

Not we construct f by smoothing the length function as follows. Fix a point $x_0 \in M$. Let $f_0 : M \to \mathbb{R}$ be the length from x_0. Then f_0 is a Lipschitz continuous function with Lipschitz constant 1. Since M is complete, f_0 is a proper function such that $f_0^{-1}((-\infty, c])$ is compact for any c. Let $\{int(D_{x_\gamma}(R_\gamma))\}$ be a locally finite open covering of M by open disks centered in x_γ with radius R_γ. Fix an isometry $TM_{x_\gamma} \cong \mathbb{R}^n$. We also assume that the exponential map centered in x_γ gives a coordinate of $M_\gamma = int(D_{x_\gamma}(R_\gamma))$, and the derivative of the exponential map and its inverse at any point has bounded by 2 with respect to operator norm. In particular f_0 has Lipschitz constant 2 for the standard metric on $\mathbb{R}^n$. We use this coordinate in the following local construction. Let $\{\rho_\gamma\}$ be a smooth partition of unity for it. Let $0 < r_\gamma < R_\gamma$ be the radius of the smallest disk centered in x_r containing the image of the support of ρ_γ. Let C_γ be the maximal value of $|d\rho_\gamma|$ for the standard metric on $\mathbb{R}^n$. Let n_γ be the number of open disks in the locally finite covering which intersects $D_{x_\gamma}(r_\gamma)$. Take a smooth function $K : \mathbb{R}^n \to \mathbb{R}$ satisfying $\int K(y)dy = 1$ and $K(y) = 0$ if $|y| > min\{1, (R_\gamma - r_\gamma)/2, 1/(n_\gamma C_\gamma)\}$. Then the smoothing of f defined by $f_\gamma(x) = \int K(x-y)f_0(y)dy$ $(x \in D_{x_\gamma}(r_\gamma))$ is Lipschitz continuous with Lipschitz constant 2 for the standard metric on $\mathbb{R}^n$, and satisfies $|f_\gamma(x) - f_0(x)| < min\{1, 2/(n_\gamma C_\gamma)\}$ for $x \in D_{x_\gamma}(r_\gamma)$. Now define f to be $\sum_\gamma \rho_\gamma f_\gamma$. Then $|f - f_0| \leq 1$. In particular f is also a proper map and $f^{-1}((-\infty, c])$ is compact for any c. Decompose df as follows: $df = (\sum_\gamma \rho_\gamma df_\gamma) + (\sum_\gamma (d\rho_\gamma) f_0) + (\sum_\gamma d\rho_\gamma (f_\gamma - f_0))$ Since the second term is zero, we have $|df| \leq \sum_\gamma \rho_\gamma \cdot |df_\gamma| + \sum_\gamma |d\rho_\gamma||f_\gamma - f_0|$. Both terms are bounded from our construction. ☐

References

1. M. F. Atiyah and I. M. Singer, *The index of elliptic operators. I*, Ann. of Math. (2) **87** (1968), 484–530.
2. P. R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, J. Functional Analysis **12** (1973), 401–414.
3. V. Danilov, *The geometry of toric varieties (Russian)*, Uspekhi Mat. Nauk **33** (1978), no. 2, 85–134, English translation: Russian Math. Surveys 33 (1978), no. 2, 97–154.
4. H. Fujita, M. Furuta, and T. Yoshida, *Torus fibrations and localization of index I -Polarization and acyclic fibrations*, to appear in J. Math. Sci. Univ. Tokyo. Also available at arXiv:0804.3258.
5. ———, *Torus fibrations and localization of index III -Equivariant version and its applications*, in preparation.
6. M. Furuta, *Index theorem 1*, Translations of Mathematical Monographs, vol. 235, American Mathematical Society, Providence, RI, 2007, Translated from the 1999 Japanese original by Kaoru Ono, Iwanami Series in Modern Mathematics.
7. R. E. Greene and H. Wu, *C^∞ approximations of convex, subharmonic, and plurisubharmonic functions*, Ann. Sci. École Norm. Sup. (4) **12** (1979), no. 1, 47–84.
8. M. Gromov, *Kähler hyperbolicity and L^2-Hodge theory*, J. Differential Geom. **33** (1991), no. 1, 263–292.
9. M. D. Hamilton, *Locally toric manifolds and singular Bohr-Sommerfeld leaves*, arXiv:0709.4058.
10. H. B. Lawson and M. L. Michelsohn, *Spin geometry*, Princeton Math. Series, vol. 89, Princeton University Press, Princeton, NJ, 1989.
11. M. Symington, *Four dimensions from two in symplectic topology*, Topology and geometry of manifolds (Athens, GA, 2001), Proc. Sympos. Pure Math., vol. 71, Amer. Math. Soc., Providence, RI, 2003, pp. 153–208.
12. J. Williamson, *On the algebraic problem concerning the normal forms of linear dynamical systems*, Amer. J. Math. **58** (1936), no. 1, 141–163.
13. T. Yoshida, *Local torus actions modeled on the standard represenation*, arXiv:0710.2166, 2007.
14. ———, *On liftings of local torus actions to fiber bundles*, Toric Topology, Contemp. Math., vol. 460, Amer. Math. Soc., Providence, RI, 2008, pp. 391–401.

DEPARTMENT OF MATHEMATICS, GAKUSHUIN UNIVERSITY, 1-5-1 MEJIRO, TOSHIMA-KU, TOKYO, 171-8588, JAPAN
E-mail address: `hajime@math.gakushuin.ac.jp`

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan
E-mail address: `furuta@ms.u-tokyo.ac.jp`

Department of Mathematics, Graduate School of Science and Technology, Meiji University, 1-1-1 Higashimita, Tama-ku, Kawasaki, 214-8571, Japan
E-mail address: `takahiko@math.meiji.ac.jp`